

Pulmonary and Generalized Lysosomal Storage Induced by Amphiphilic Drugs

by Zdenek Hruban*

Administration of amphiphilic drugs to experimental animals causes formation of myelinoid bodies in many cell types, accumulation of foamy macrophages in pulmonary alveoli and pulmonary alveolar proteinosis. These changes are the result of an interaction between the drugs and phospholipids which leads to an alteration in physicochemical properties of the phospholipids. Impairment of the digestion of altered pulmonary secretions in phagosomes of macrophages results in accumulation of foam cells in pulmonary alveoli. Impairment of the metabolism of altered phospholipids removed by autophagy induces an accumulation of myelinoid bodies. The administration of amphiphilic compounds thus causes pulmonary intra-alveolar histiocytosis which is a part of a drug-induced lysosomal storage or generalized lipidosis.

The accumulation of drug-lipid complexes in myelinoid bodies and in pulmonary foam cells may lead to alteration of cellular functioning and to clinical disease. Currently over 50 amphiphilic drugs are known. Unique pharmacological properties necessitate clinical use of some of these drugs. The occurrence and severity of potential clinical side effects depend on the nature of each drug, dosage and duration of treatment, simultaneous administration of other drugs and foods, individual metabolic pattern of the patient and other factors. Further studies on factors preventing and potentiating adverse effects of amphiphilic drugs are indicated.

In recent years it has become apparent that certain drugs administered systematically to man have serious side effects because of their affinity for the lungs. The pulmonary pathology produced by these drugs has received relatively little clinical attention because of the insidious and chronic nature of its development. With increasing awareness that a large variety of seemingly harmless drugs can induce such lung changes, it is imperative that more rigorous studies be carried out before such drugs are put to clinical use. The purpose of this report is to review a group of drugs which lead to intra-alveolar histiocytosis and to generalized lysosomal storage.

Foam Cells

In 1966, Greselin reported (1) that chronic administration of a drug which inhibits cholesterol synthesis, *trans*-1,4-bis(2-chlorobenzylaminomethyl)cyclohexane dichloride, also called AY-9944, induces accumulation of foam cells in pulmonary alveoli of experimental animals. In the past 15 years other drugs with a variety of pharmacological effects were shown to produce similar pulmonary side effects (Table 1).

These drugs induce essentially similar histological and ultrastructural changes in the lung, although the

time required for the appearance of the foam cells and their quantity and size may be quite different in various animal species (1-4). The lungs of the treated animals increase in weight (1,5) and contain whitish plaques and nodules on gross examination (1,6). Microscopically these plaques consist of intra-alveolar accumulations of foam cells (1,7).

The foam cells are between 20 and 80 μ m in diameter (4,8-10) and have a centrally placed nucleus. The cytoplasm is abundant, pale, and finely reticulated in hematoxylin-eosin stained sections (Fig. 1). In toluidine blue-stained Epon sections, the foam cells are filled with numerous round inclusions. Staining of foam cells with Sudan Black B and with Baker's acid hematein for phospholipids are positive, while staining with Sudan III and by the periodic acid-Schiff method are usually negative (5,7,10,11). Examination of fresh tissues in polarized light identifies both the inclusions in foam cells and myelinoid bodies as Maltese cross figures (11,12). Histochemical studies reveal high activities of acid phosphatase and β -glucuronidase, implicating the foam cells as macrophages (4,10,13,14).

The time sequence of pulmonary changes associated with foam cell accumulation has been studied at light and electron microscopic levels. The first change observed in iprindole-treated rats is interstitial pulmonary edema associated with degenerative changes in capillary endothelia (14). Subsequently endothelia and alveolar epithelia become swollen and the septa infil-

*Department of Pathology, University of Chicago, Chicago, IL 60637.

Table 1. Inducers of intra-alveolar foam cells.^a

Alveolar lumen			Type II cells		Myelinoid bodies			Chlorphen- termine binding	Affinity for lung	Other effects	References
Secretions	Foam cells	Protein- osis	Hyper- trophy	Hyper- plasia	Type I cells	Clara cells	Other cells				
Amiodarone	+						+				(243)
Amiripryline	0				+	+		+			(3,41,118,244)
AY-9944	+		+					+			(1,8,144)
Boxidine	+										(26,7)
Chlorcyclizine	+		+		+	+	+	+			(7,26,118,221,245)
1-Chloroamiripryline	+		+		+		+				(3,41)
1-Chloro-10,11-dehydro- amiripryline	+		+		+		+				(3,41)
Chloroquine	+		+		+	+		+			(7,26,118)
Chlorphentermine	+	+	+	+	+	+	+	+		A	(3,10,19,21,114,118, 145,245,248,249,250)
Chlorpromazine	0										(3,41,244,245)
Clofex	+							+			(5,17,246)
Clomipramine	+				+		+	+			(3,41,113)
Cycizine	+		+					+			(7,26,221,245)
4,4'-Diethylamino- ethoxyhexestrol	+				+		+				(33,205)
Erythronycin	+	+						+			(212)
Ethyl fluclozepate	+										(247)
Fenfluramine	+										(2)
Fluoxetine	+										(24)
Haloperidol	+										(7,26)
Homochlorcyclizine	+										(7,26)
Hydroxyzine	+										(7,26)
Imipramine	+		+		+		+	+			(3,41,118,221,244,245,251)
Iprindole	+	+			+	+	+				(3,14,16,20,41)
Medizine	+										(7,26)
Norchlorcyclizine	+										(7,26,112)
Noxiptiline	0										(3,41)
Perhexiline	+							+			(123)
Penthermine	0										(145,248)
Quinacrine	+							+		A	(245)
RMI 10.393	+										(11)
Thioridazine	0										(3,41)
Triparanol	+		+			+	+				(7,9,26)
Zimeldine	+						+				(243)

^aLegend: +, present; 0, absent; A, induces pulmonary hypertension in man (206); B, induces cytochrome P-450 (221); C, metabolized in lung (221).

trated by interstitial macrophages (14). The lungs of chlorphentermine-treated rats show hyperemia, aggregation of leucocytes in venules and perivascular infiltration by monocytes during the first week of treatment (4). In general, a few intra-alveolar macrophages appear early and become progressively larger, binucleated, and more numerous (5,7,10,15). These macrophages are derived from interstitial macrophages (10,16), which in turn originate from blood monocytes (4,17). In histological sections the alveolar lumina contain abundant amorphous material. On ultrastructural examination the material is identified as secretions derived from secretory vacuoles of granular pneumocytes (Figs. 2 and 3). It has been shown (7,18,19) that the intra-alveolar macrophages phagocytize the secretions, which are rich in dipalmitoyllecithin (9), and become foam cells (Fig. 4). After 3–6 weeks of treatment, the foam cells are numerous (4,7,8,16).

A marked decrease in the secretory activity of Type II pneumocytes is observed after 9 months of treatment (20). After 12 months of iprindole feeding, the accumulations of foam cells are replaced by pale eosinophilic granular material in the intra-alveolar spaces as a result of cellular breakdown, and the histological picture is that of alveolar proteinosis (20). Chronic administration of chlorphentermine and of AY-9944 will lead to similar changes (1,21). If the administration of the drug is withdrawn after several weeks of treatment, the size, the number of foam cells, the ultrastructure and enzyme activities will normalize in 2–5 weeks (7,10,22–24). During this process the secretions in phagosomes of macrophages are replaced by electron dense heterogeneous material (7). The current knowledge of the alveolar macrophage in drug-induced lipidoses has been reviewed by Reasor (23).




FIGURE 1. Densely packed foam cells in alveolar lumina of a rat fed chlorcyclizine. Hematoxylin-eosin, $\times 590$.




FIGURE 3. Secretion of a granular pneumocyte from a rat treated with chlorcyclizine (302). $\times 5600$.




FIGURE 2. Portion of an alveolus filled with concentric and tubular secretions from a rat treated with chlorcyclizine. $\times 6900$.




FIGURE 4. Foamy alveolar macrophage with phagocytized secretions from a rat treated with chlorcyclizine. $\times 4000$.

Myelinoid Bodies

Several pulmonary cells show striking changes following treatment with amphiphilic drugs (AP drugs). Type II pneumocytes become hypertrophic and hyperplastic after triparanol and chlorcyclizine treatment (7,8) but do not show changes after chlorphentermine treatment (10). The secretory vacuoles of Type II pneumocytes become enlarged (7,14). Myelinoid bodies (25,26) and heterogeneous dense bodies appear in the cytoplasm of Type I pneumocytes (Fig. 5), ciliated bronchiolar epithelia and Clara cells, smooth muscle cells, fibroblasts and capillary endothelia (Table 1).

Drug-induced myelinoid bodies (DIM bodies) are secondary lysosomes containing lamellar, reticular or crystalloid material (Figs. 6 and 7). The lamellae form

concentric layers with a 40 to 45 Å periodicity (27). Reticular or crystalloid myelinoid bodies are found in various pulmonary cells, but not in pulmonary macrophages (90). Heterogeneous dense bodies are secondary lysosomes filled with heterogeneous material.

Myelinoid bodies have received a variety of names. The original term myeloid body (25,26,28) should be, in the opinion of Ghadially (29), replaced by the more appropriate term myelinoid body. Other investigators prefer to use terms such as body, inclusion or lysosome preceded by adjectives such as lamellar, lamellate, multilamellate, honeycomblake and membranous. Names which do not differentiate between membrane-bound myelinoid bodies and myelin figures without limiting membranes should be avoided.

The drugs which induce accumulations of pulmonary




FIGURE 5. Type I pneumocytes from a rat fed chlorcyclizine for 5 weeks. The upper cell contains numerous myelinoid bodies. The lower cell contains heterogeneous dense bodies and material resembling alveolar secretions. $\times 15,600$.




FIGURE 6. Myelinoid bodies induced in a rat hepatocyte by chlorcyclizine. The upper body is of the reticular type, the lower body is of the mixed type. $\times 23,400$.

foam cells also induce the formation of myelinoid bodies in many other cell types of extrapulmonary tissues (Fig. 8) both *in vivo* (Table 2) and *in vitro* (Table 3). High doses of chlorphentermine, chloramitryptiline, RM1 10.393, and erythromycin induce formation of foam cells at extrapulmonary sites particularly in lymph nodes, spleen and thymus (11,30). Patients treated with diethylaminoethoxyhexestrol (DEAEH) have foamy cells with deep blue granules in their bone marrow (31,32), spleen and lymph nodes (33). Foam cells do not, however, appear in similarly treated rats (34).

Because the inclusions in foam cells and the DIM-bodies are modified lysosomes, knowledge of lysosome formation by heterophagy and by autophagy is necessary to understand their nature (26,35). In autophagy, sequestering cisternae surround a portion of the cytoplasm containing organelles and form an autophagic vacuole. The thin membranes of the sequestering cisterna are then transformed into a thick limiting




FIGURE 7. Crystalloid type of a myelinoid body from the corpus luteum of a chlorcyclizine-treated rat. $\times 22,400$.

membrane of the lysosome. Hydrolytic enzymes formed in endoplasmic reticulum are brought to the autophagic vacuole within primary lysosomes. The sequestered organelles are broken down by the hydrolytic enzymes and the autophagic vacuole changes into a heterogeneous dense body (secondary lysosome). Further digestion leads to the appearance of smaller homogeneous bodies (residual bodies). In heterophagy, the extracellular material enters the cell within pinocytotic vesicles and phagocytic vacuoles. Primary lysosomes empty their enzymes into this vacuole, and the digestion proceeds as in autophagy.

If the degradation of substrates sequestered within lysosomes is impaired, the substrates will accumulate, and the lysosomes will become storage bodies. Lysosomal digestion may be impaired by several mechanisms (36,37). For example, each of the human storage diseases is caused by the absence of a specific lysosomal enzyme. Myelinoid bodies are storage bodies containing lipids of membranes whose digestion is impaired by drugs (18). The inclusions in foam cells are phagosomes containing phagocytized pulmonary secretions whose digestion is impaired by drugs.

The inclusions in foam cells (Fig. 4) and the DIM-bodies (Fig. 6) should be differentiated from secretory granules of Type II pneumocytes (38) (Fig. 3). All three structures are limited by a 90 Å thick membrane and contain lamellar material. The lamellae within phagosomes of pulmonary macrophages are less densely packed and less orderly arranged than those of DIM-bodies. Tangential cuts through myelinoid bodies can obscure the outlines of the limiting membrane, leading

Table 2. Inducers of myelinoid bodies *in vivo*.^a

Drugs and chemicals: name, generic (brand) ^b	Therapeutic use ^c	Amphiphilic molecule	Myelinoid bodies <i>in vitro</i>	Myelinoid bodies in animal	Myelinoid bodies in man	Adverse effects in man	Alveolar foam cells ^d	Liver	Kidney	Central NS ^e	Peripheral NS ^e	Pigment epithelium, retina ^f	Müller cells ^f	Neurons of retina ^f	Visual cells ^f	Lens ^g	Cornea ^h	Lymphatic system ⁱ	Blood ^j	Striated muscle ^k	Adrenal	Testis	Fetus	Malformation, abortion	References
(2-N-Methylpiperazinomethyl-1,3-diazofluoro-anthen-1-oxide (AC-3579)	X	+	+	+	+	+	+	+	+	0	+	+	0	+	0	+	+	+	+	+	+				(60,62)
Amiodarone (Cordarone)	E	+	+	+	+	+	+	0	0	0	+	+	0	+	0	+	+	+	+	+	+				(243,252)
Amiripryline (Elavil)	H	+	+	+	+	+	+	0	0	0	+	+	+	+	+	+	+	+	+	+	+				(3)
1-Chloroamiripryline, R044318	H	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(132,83,84,142)
1-Chloro-10,11-dehydroamiripryline	H	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(3)
<i>trans</i> -1,4-bis(2-Chlorobenzylaminoethyl)-cyclohexane (AY 9944)	T	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(8,77,144,267)
Azasterol: 20,25 diazcholesterol	T	0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(77,253)
Bilirubin	T	+	+	+	+	+	+	0	+	+	+	+	+	+	+	+	+	+	+	+	+				(254,255)
Boxidine	T	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(7)
Chloroquine (Aralen)	M	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(6,7,26,58,202,268,269)
Chlorpheniramine (Pre-sate)	C	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(10,13,41,132)
Phentermine (Phentermine)	C	+	+	+	+	+	0	+	+	+	+	+	+	+	+	+	+	+	+	0	+				(145,256)
Chlorpromazine (Thorazine)	V	+	+	+	+	+	0	0	0	+	+	+	+	+	+	+	+	+	+	0	+				(8,188,270)
Clindamycin (Cleocin)	F	0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(212,30)
Clofex	C	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(17)
Clozapine (Anafranil)	H	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(3,41)
Cyanochlorophenylamidinourea	VX	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(257)
Cyclizine (Marezine)	M	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(41)
Cyclochlorizine (Fedrazil)	JLT	+	+	+	+	+	+	+	+	+	+	+	+	+	0	+	+	+	+	+	+				(86,7)
Chlorcyclizine	LT	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(7,26,271)
Homochlorcyclizine	LTY	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(7,97)
Hydroxyzine (Atarax)	TX	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(7)
Meclizine (Bonine)	JL	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(7)
Norchlorcyclizine	T	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(7)
4,4'-Diethylaminoethoxyhexestrol	E	+	+	+	+	+	+	+	+	0	+	+	+	+	+	+	+	+	+	+	+				(28,31,33,34,96,103,205)
Erythronycin (Ilotycin)	F	0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(212,30)
Ethyl fluclozepate	H	0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(247)
Fenfluramine (Pondimin)	C	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(41)
Fluoxetine, [<i>N</i> -methyl-3-phenyl(α,α -trifluoro- <i>p</i> -tolyl)oxypropylamine hydrochloride]	Y	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(24)
Gentamycin (Garamycin)	F	0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(92,183,210,272)
Haloperidol (Haldol)	OX	0	+	+	+	+	+	0	+	+	+	+	+	+	+	+	+	+	+	+	+				(7)
6-Hydroxydopamine, 5-hydroxydopamine	W	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(211)
IA3, 8-chloro-5-methyl-2-[2-(diethylamino)-ethyl]-2H-(1-benzothio)pyranone[4,3,2-c,d]indazole	W	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(257)
Imipramine (Tofranil)	H	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(3,41)
Indoramine	A	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(257)
Ipindole (Prondol)	H	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(3,14,16,20)
Maprotiline (Ludomil)	H	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(101,102)
Nortriptyline (Aventyl)	H	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				(258)

Table 2. Inducers of myeloid bodies *in vivo*.^a (continued)

Drugs and chemicals: name, generic (brand) ^b	Therapeutic use ^c	Amphiphilic molecule	Myeloid bodies <i>in vitro</i>	Myeloid bodies in animal	Myeloid bodies in man	Adverse effects in man	Alveolar foam cells ^d	Liver	Kidney	Central NS ^e	Peripheral NS ^e	Pigment epithelium, retina ^f	Müller cells ^f	Neurons of retina ^f	Visual cells ^f	Lens ^g	Cornea ^h	Lymphatic system ⁱ	Blood ^j	Striated muscle ^k	Adrenal	Testis	Fetus	Malformation, abortion	References
Noxiptiline (Agedal)	H	+	+	+	+	+	0	0	+	0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(222) (185) (40,61,73,87,123-125, 273-275)
Paraquat	S	0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(273-275)
Perhexiline (Pexid)	ET	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(42) (97) (87,259,12) (27)
Phenacetin, Acetophenetidine	B	0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(132,84)
Preynilamine	E	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(260)
Quinacrine	M	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(41,303) (222)
R-800, cis-7-fluoro-1-phenyl-3-isochroman- methylamine	C	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(175,261,262,304) (25,35,77,78,276-279)
RMI 10,393, 5-[p-(fluoren-9-ylidenemethyl)- phenyl]-2-piperidineethanol	G	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(243,252)
SKF 14336-D, 9-(3-dimethylaminopropyl)-2- chloroacridane	X	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(260)
Tamoxifen (Nolvadex)	K	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(41,303) (222)
Thioridazine (Mellaril)	VX	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(175,261,262,304) (25,35,77,78,276-279)
Tilorone	F	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(243,252)
Triparanol (MER-29)	T	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(243,252)
Zimelidine	H	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	(243,252)

^a Legend: +, present; ++, abundant; ±, infrequent; 0, absent.^b Additional brand names are found in The Merck Index.^c Uses: A, adrenergic blocker, sympatholytic; B, analgesic; C, anorectic; D, antagonist of serotonin; E, antianxiety; F, antibacterial, antiviral; G, anticholinergic; H, antidepressant; I, antidote; J, antiepileptic, antinauseant; K, antestrogenic; L, antihistaminic; M, antimalarial, antihelminthic; N, antipruritic; O, antipsychotic; P, dye; Q, estrogenic; R, gonadal stimulating, ovulatory; S, herbicide; T, hypocholeraemic; U, myocardial depressant; V, neuroleptic; W, schistosomicide; X, sedative, tranquilizer; Y, serotonin antagonist; Z, suppressant of lupus erythematosus, antirheumatic.^d See Table 1.^e Ref. (263).^f Refs. (126,178,216,264).^g Refs. (39,201).^h Ref. (121).ⁱ Ref. (41).^j Refs. (32,41,99,265).^k Ref. (266).

Table 3. Inducers of myelinoid bodies *in vitro* systems.^a

Drugs and chemicals: name, generic (brand) ^b	Thera- peutic use ^c	Amphi- philic molecule	Culture of nervous tissue	Culture of macrophages	Culture of other cells	Other systems	References
Acridine orange					+		(280)
1-Amino-4-octylpiperazine, AP22	F	+	+				(281)
1,7-bis(<i>p</i> -Aminophenoxy)heptane, 153C51	W	+				d,e	(282,108)
Amitriptyline	H	+		+			(283)
AY 9944	T	+	+				(215)
Benzylidene-1-aminopiperazine, AP8	F	+	+				(281)
Bilirubin		0	+				(284,285)
Brompheniramine	L	+		+			(257)
Chloroquine	M	+	+	+	+	f	(107,290-292)
Chlorpheniramine (Teldrin)	L	+		+			(257)
Chlorpromazine (Thorazine)	V	+	+	+		g	(188,283,293)
Clociguanil	M	+			+		(41)
Diethylaminoethoxyhexestrol	E	+			+		(286)
Gentamycin	F	0			+		(210)
6-Hydroxydopamine		0			+		(211)
Lysergide	HXY	+	+				(172)
Mesoridazine (Serentil)	OX	+	+				(258)
Mianserine (Tolvin)	LNy	+			+		(257)
Myoinositol		0	+				(287)
Nortriptyline	H	+	+				(258)
Perhexiline	ET	+	+		+		(73,294,295)
Pheniramine	L	+		+			(257)
Phentermine	C	+		+			(283)
Promethazine (Phenergan)	JL	+	+				(288)
Trifluoperazine (Stelazine)	OX	+	+				(288)
Trimeprazine (Temaril)	LN	+	+				(288)
Triparanol (MER-29)	T	+			+		(100)
Tunicamycin	F				+		(289)

^a Legend: +, present; 0, absent.

^b Additional brand names are found in The Merck Index.

^c See Table 2.

^d Schistosomes contain myelinoid bodies.

^e Retinotoxic.

^f Lysosomes of malarial parasites altered, no myelinoid bodies.

^g In perfused rat liver.

to the misinterpretation that they are lacking a limiting membrane (39-42).

Drug-induced myelinoid bodies should be clearly differentiated from other cellular structures containing concentrically arranged lamellae (26). Structures simulating DIM bodies can be grouped into categories of cytoplasmic organelles, secretory granules (43,44), pigmented granules of neurons (45), storage granules (46), autophagic vacuoles in cells rich in phospholipids (47), phagocytic vacuoles oversupplied with phospholipids (42), and secondary lysosomes with a slow phospholipid breakdown (48,49). The term myelinoid bodies should not be applied to secondary lysosomes or vacuoles containing a few membrane fragments or an occasional myelin figure.

DIM bodies show a superficial resemblance to smooth fingerprints (50,51) and to whorls formed by smooth portions of rough cisternae (52), both of which may be also induced by amphiphilic drugs. Myelinoid bodies should be distinguished from myelin figures lying free in the cytoplasm or on the surface of organelles (26,53). Myelin figures represent bilayered lamellae of amphi-

philic lipids which may arise from altered cellular membranes, in neutral lipids during lipolysis (54,55) or from altered myelin (56). Most often they are artifacts seen in tissue fixed in aldehydes followed by osmium tetroxide (57). Extracellular and intraluminal myelin figures may represent extruded contents of myelinoid bodies (Fig. 8) (42,58,59).

Biochemical Studies

Since the establishment of the lipid nature of the material stored within the foam cells and within myelinoid bodies, various investigators have analyzed the biochemical composition of organs (60-62), tumor cells (63), isolated alveolar macrophages and isolated DIM bodies derived from animals and people treated with amphiphilic drugs (AP drugs). As reviewed by Lüllmann-Rauch (41), the tissues show a marked increase of total phospholipids (PLs) and variable increases of cholesterol, triglycerides and gangliosides. The absolute increase in PLs is usually due to an increase of phosphatidylcholine and in decreasing per-




FIGURE 8. Renal collecting tubule from a rat treated with chloroquine for 6 weeks. The cells contain numerous myelinoid bodies. The lumen contains cellular debris with numerous myelinoid bodies and a few mitochondria derived from ruptured proximal cells. $\times 4,000$.

centages to phosphatidylinositol, phosphatidylethanolamine, bis(monoacylglycerol)phosphate (BMGP), sphingomyelin and phosphatidylserine.

Chloroquine and DEAEH increase the incorporation of glycerol into lysosomal phospholipids, but decrease its incorporation into microsomal phospholipids. These drugs greatly increase the synthesis of phosphatidylglycerol in rat liver (64). Homochlorcyclizine induces accumulation of glycolipids, phosphatidylcholine and BMGP in the liver (65). The chlorphentermine-induced accumulation of glycoproteins and mucopolysaccharides in culture of intestinal cells is interpreted to be the consequence of crinophagy rather than endocytosis (66).

Excessive storage of gangliosides is demonstrable in miniature pigs treated with chloroquine. In the brain,

GM₂ ganglioside increases while phospholipids and cholesterol are unchanged (67). Skeletal muscle GM₂ ganglioside increases 100-fold, BMGP 50-fold and phospholipids 3-fold (68). Liver, spleen and lungs show a marked increase of glucosylceramide, GM₂ ganglioside and BMGP (69). Chlorphentermine induces a 14% increase of GM₁ ganglioside in rat brain (70). The incorporation of phosphorus into phosphatidylcholine is reduced and into phosphatidic acid increased by several amphiphilic drugs studied in cerebral cortex mince (71). Perhexiline induces the accumulation of gangliosides in human liver (72) and peripheral nerve (73) as well as that of GM₃ ganglioside in cultured fibroblasts (74).

The composition of DIM bodies and of tissues containing these bodies depends on the type of tissue examined, the animal species, diet (75), duration of treatment and

the type of the drug. The excess of hepatic phospholipids can be explained by the increase of PLs in myelinoid bodies alone (76). Several investigators studied the composition of cellular fractions rich in DIM bodies (33,77,78). The composition of smooth microsomal membranes from livers of rats treated with AC 3579 for 4 days is altered. The phospholipid/protein ratio is markedly increased while the cholesterol/phospholipid ratio is decreased. The isolated DIM bodies have a similar composition, suggesting that the content of myelinoid bodies could be derived from sequestered membranes of smooth endoplasmic reticulum. The labeled drug is found to be selectively concentrated in DIM bodies (79). The effect of this drug on membrane microviscosity is cholesterol-like (80). The low cholesterol/phospholipid ratio in DIM bodies could be the result of replacement of free cholesterol in membranes by the amphiphilic drugs (76).

Biochemical studies performed on whole lungs or on isolated alveolar macrophages obtained from animals treated with AP drugs reveal alterations in the content and composition of lipids. Cloforex increases the cholesterol content of rat lung by 50% and the phospholipid content five times (5). Perhexiline, however, decreases the concentration of cholesterol, PLs and gangliosides in rat lungs (61). Alterations induced in the lungs by chlorphentermine were reported by several investigators. Seiler and Wasserman (81) demonstrated a marked increase of phosphatidylcholine, and an increase in sphingomyelin, free cholesterol and cholesterol ester. Other investigators observed an increase in total PLs, phosphatidylcholine, phosphatidylinositol, phosphatidylglycerol, sphingomyelin, free cholesterol and triglycerides (11,82-84). The fatty acids of phospholipids were more saturated, the content of phosphatidylethanolamine was decreased (82) and the surface tension forces were reduced (85). RMI 10.393 produces changes similar to those seen with chlorphentermine (86). An increase of PLs in the lung lavage fluid occurs early (87).

The alveolar macrophages of rats treated with chlorphentermine and RMI 10.393 show an increase in total PLs, phosphatidylcholine, phosphatidylinositol, phosphatidylglycerol, phosphatidic acid, sphingomyelin and free cholesterol (11,83,84,88,89). 1-Chloramitryptiline, on the other hand, leads to a slight accumulation of lecithin and of phosphatidic acid in the lung, while the foam cells accumulate cholesterol, cholesterol esters and free fatty acids. The authors conclude that chlorphentermine and RMI 10.393 impair secretory capacity of pneumocytes, while chloramitryptiline does not. The alveolar macrophages induced by chlorphentermine show an increased lipid peroxidation, probably occurring within lysosomes, as well as augmented antioxidant defense mechanisms (54,90).

A marked increase of the levels of total PLs, total sterols, BMGP and desmosterol is demonstrable in the lamellar body fraction isolated from lungs of rats treated with DEAEH (33). Desmosterol is also found in fractions containing myelinoid bodies induced by inhibitors of cholesterol synthesis (91,92).

Tissues of animals and patients treated with various amphiphilic drugs contain an unusual phospholipid, bis(monoacylglycero)phosphate, or BMGP (33,93-97). It forms a much greater proportion of phospholipids in human tissues than in tissues of animals (41,96), probably because of a difference in drug metabolism (98).

The presence of lysosomal enzymes within DIM bodies in cells of different organs and species, after the administration of various AP drugs, has been repeatedly demonstrated (25,32,41,99-102). Quantitative studies usually reveal increased activities of acid phosphatase, β -glucuronidase, β -galactosidase, and decreased activities of acid esterase, β -*N*-acetylglucosaminidase and cathepsin D (62,72,88,103-108). The levels of several lysosomal enzymes (particularly that of β -*N*-acetylglucosaminidase) of foam cells are increased, while acid esterase and α -mannosidase are decreased (23,88,109,110). Chlorcyclizine inhibits the supernatant enzyme steryl ester hydrolase *in vitro* (111). Myelinoid bodies of chloroquine and DEAEH-treated rats accumulate cholesterol ester because these drugs block its lysosomal degradation (76).

The accumulation of PLs within tissues, DIM bodies and foam cells has been correlated with the accumulation of amphiphilic drugs and of their metabolites. Several AP drugs and/or their metabolites are localized mostly in the lung and spleen, and moderately in the liver, kidney, heart, muscle, adrenals and fat (112-116). The number of DIM bodies appears to be related to the concentration of the drug in a tissue. Correlation between the concentration of acidic PLs and accumulation of the drug and its metabolites suggesting formation of a stable complex is reported by Matsuzawa et al. (98,117) in livers of patients treated with DEAEH. The binding of chlorphentermine to phospholipid-rich subcellular fractions is inhibited by desmethylinipramine and other basic amines (118) (Table 1). Several investigators demonstrated the accumulation of AP drugs within isolated myelinoid bodies (76,78,119). In addition to their accumulation within DIM bodies, some AP drugs form reversible complexes with phospholipids of the smooth endoplasmic reticulum (120). Such binding may decrease the activity of microsomal enzymes.

Preferential Organ Involvement

The number of DIM bodies in a tissue will depend on the speed with which a given AP drug is metabolized, the duration of treatment and the dosage of the drug (121). Potent AP drugs can induce DIM bodies in all tissues, but the quantity of DIM bodies will depend on the cell type. Weak AP drugs induce DIM bodies in a few cell types only, which usually include the lymphocytes. The sensitivity of lymphocytes may be related to the rapid turnover of glycerolipids in these cells (122). Perhexiline exemplifies a drug which induces DIM bodies in an unusual distribution pattern: Myelinoid bodies are commonly found in the retina, dorsal root ganglia, adrenal gland and muscle cells of the rat, while they are few in the hepatocytes and lymphocytes, and

foam cells in the lung are scarce (123). The same drug is, however, associated with the cirrhosis of the liver in man (124,125).

The difference in the amounts of DIM bodies in specific organs after treatment with individual AP drugs is probably related to the lipid composition of the tissue, binding of the drug by the tissue, metabolism and excretion of the drug. Certain cell types may contain specific phospholipids for which certain AP drugs have selective affinities. Thus the absence or presence of a specific phospholipid may decide whether an AP drug will induce myelinoid bodies in one or another cell type in a given tissue, such as a neuron or pigment epithelium of the retina (126). The binding of AP drugs to liposomes made of different lipids occurs in this increasing order: sphingomyelin and phosphatidylcholine, phosphatidylethanolamine, and phosphatidylserine (127). The divalent chloroquine has a very high affinity for phosphatidylserine and gangliosides which may explain its tendency to affect ganglionic cells. The positively charged adriamycin shows a high affinity for cardiolipin, which may be related to its selective toxicity for mitochondria (128) and its failure to induce typical myelinoid bodies.

The involvement of the central nervous system depends on the efficiency of the blood-brain barrier (87,129). The accumulation of AP bodies in liver cells may be facilitated by a slow metabolic inactivation of the drug, fast metabolism of the drug into an active metabolite and/or by passage of the drug through the hepatocytes during excretion of the drug in bile. Several AP drugs are excreted as bile salt-lecithin-drug micelles (130). The lungs also contain drug-metabolizing enzymes (131,132).

Uptake of Amphiphilic Drugs by the Lungs

The uptake of AP drugs has been studied in isolated perfused lungs of rabbits and rats. The accumulation has a saturable component and a linear nonsaturable component. The saturable component is attributed to specific binding sites or to a carrier-mediated transport. The linear component is attributed to diffusion to intracellular and extracellular fluid and to nonspecific binding to cellular components. The uptake of the drug is directly related to the degree of protonation of the nitrogen atom and the accumulation is reversible (133). Minchin et al. (134) suggested that the transport saturability could be explained either as a carrier-mediated transport or as an alteration of diffusion because of drug-induced changes in the fluid state of membranes. Uptake of chlorphentermine is a Na^+ -dependent, carrier-mediated transport process (133). Methadone forms a slowly effluxable pool in the lung comparable to chlorphentermine and may be expected to induce phospholipidosis (135). The uptake of chloroquine by pancreatic islets requires neither energy-dependent active transport nor pinocytosis (136). Lungs of chlorphentermine-pretreated rats accumulate chlorphentermine more rapidly than control lungs (137).

Pathogenesis

The storing of phospholipids and of amphiphilic drugs in both myelinoid bodies and vacuoles of alveolar foam cells suggests that the pathogenesis of both lesions is similar. The two main factors which should be considered in the pathogenesis of phospholipid accumulation within myelinoid bodies are: an increased supply of PLs to secondary lysosome through membrane or product flow (33) and a decreased removal or catabolism of the contents of secondary lysosomes. Decreased catabolism of lysosomal contents could be associated with formation of cholesterol precursors (1,7,8,28,77,78), inhibition of the fusion between primary and secondary lysosomes (58), direct inactivation of certain lysosomal enzymes (10,28,33,121) and formation of drug-lipid complexes with altered physicochemical properties (7,9,10,28,81).

The factors to be considered in the pathogenesis of the accumulation of phospholipids within intra-alveolar foam cells are similar to those just listed. The pulmonary PLs are found in the pulmonary surfactant. Accumulation of foam cells could result from an increased production of the surfactant, decreased catabolism of the surfactant within pulmonary macrophages or decreased removal of the macrophages from the lungs (8,20,26). The pulmonary surfactant has been shown to be synthesized in Type II pneumocytes (138,139). It is catabolized in alveolar macrophages (140) and in Type I and II pneumocytes (141). The phospholipids of the surfactant contain about 76% of lecithin (particularly dipalmitoylphosphatidylcholine), and 11% of phosphatidylglycerol. Lamellar inclusion bodies of Type II pneumocytes have a similar composition. Studies on the incorporation of labeled palmitic acid into pulmonary PLs showed that phospholipid storage after chlorphentermine and RMI 10.393 is the result of decreased degradation, while the slight storage of PLs induced by chloramitriptyline is the result of marked increase of synthesis in spite of an increased degradation of PLs (142).

Fetal lungs of animals treated with AP drugs contain alveolar macrophages (143) or show hyperplasia of Type II pneumocytes (144). The neonatal lung is, however, less susceptible to chlorphentermine-induced alveolar foam cell formation than is the lung of adult rats (11,23,145,146).

Drug-Lipid Interactions

The majority of drugs which induce pulmonary intra-alveolar histiocytosis and generalized formation of myelinoid bodies (generalized lipidosis) are cationic amphiphilic compounds. They contain a hydrophobic region and a primary or substituted amine group which can bear a net positive charge. Such compounds interact with anionic groups of acidic phospholipids (such as phosphatidate, phosphatidylinositol, phosphatidylserine) of membranes. The hydrophobic regions of the drugs partition into the membrane. The interaction leads to neutralization of the charges on the phosphate

groups and to the expulsion of the tightly bound divalent cations such as calcium. All this leads to changes in the movement, fusion, permeability, transport and receptor functions of membranes (147-150). The interaction between various AP drugs and phospholipids was demonstrated by means of nuclear magnetic resonance (151,152).

Several authors (41,121) have pointed out that the AP drugs exist in a protonized and in a nonprotonized form. The nonprotonized forms permeate through the plasma membrane and through lysosomal membranes. The protonized forms are bound to plasma membrane, membranes of endoplasmic reticulum and other cytomembranes. Nonprotonized molecules enter the lysosomes, where they are protonized because of the strongly acid milieu, increase intralysosomal pH, form complexes with polar lipids of lysosomal contents and are thus trapped within the lysosomes (41,121,153).

The administration of several cationic AP drugs leads to quantitative alterations of phospholipids in various tissues (84,96,97,122,154-156). In general, the tissues and lysosomes will contain increased proportions of anionic phospholipids (phosphatidate, phosphatidylinositol, phosphatidylglycerol, lysobisphosphate) and decreased proportions of triglycerides, and of cationic phospholipids phosphatidylcholine and phosphatidylethanolamine (157). The quantitative changes in PLs and the hypotriglyceridemic effects can be explained as the result of the observed inhibition of the enzymes phosphatidate phosphohydrolase (91,122,147,158) and phospholipase, and stimulation of phosphatidate cytidyltransferase (159) resulting in diversion of phospholipid synthesis from one to another pathway (147,149).

The accumulation of phospholipids (PLs) within lysosomes can be explained as the result of impaired degradation of PLs by the lysosomal phospholipase (121,122,149). The presence of the enzyme phospholipase A has been demonstrated in the plasma membrane, and in microsomal and lysosomal fractions (160). The lysosomal phospholipase cleaves both fatty acid ester linkages of lecithin and of phosphatidylethanolamine without formation of lyso-forms (161). Various *in vitro* and *in vivo* studies show that the enzyme phospholipase A (which acts on the hydrophobic group of the PL) but not phospholipase C (which acts on the hydrophilic part of the PL molecule) is inhibited by several amphiphilic drugs (91,119,120,162-164). Following the formation of a complex between AP drugs and PLs, the hydrophilic group of the drug prevents the formation of the enzyme-substrate complex, while the hydrophobic group allows penetration of the drug into the lipid layer and determines the stability of the drug-lipid complex. Thus, e.g., the hydrophobic group of AC 3579 is larger than that of AP 22, the AC 3579-lipid complex is more stable and the enzyme inactivation is more complete than are those of AP 22 (91). Both lysosomal phospholipases have been found inhibited by chloroquine, DEAEH and aminoglycoside antibiotics (165,166). Depression of lysosomal phospho-

lipase activities may be also due to the drug-induced increase of intralysosomal pH (153).

The above alterations in enzymes do not explain the lysosomal accumulation of the anionic phospholipid bis(monoacylglycerol)phosphate (BMGP) also called lysobisphosphatidic acid. BMGP contains very high amounts of polyenoic fatty acids, particularly docosa-hexaenoic acid (167,168). It is synthesized in lysosomes (169,170) by the BMGP synthetase by transfer of acyl groups from phosphatidylinositol to phosphatidylglycerol or lysophosphatidylglycerol (76). BMGP is resistant to degradation by lysosomal hydrolases (76). It is a normal component of lysosomal PLs, where it may account for 6 to 28% of total PLs (76,167,171). Some of the lysosomal BMGP could have its origin in the plasma membrane (167).

The lipid composition of myelinoid bodies and consequently also that of tissues is thus determined in part by the type of lipid reaching the lysosomes either by heterophagy or by autophagy. The formation of myelinoid bodies induced by lysergide has been attributed to increased endocytosis (172) which supplies lipids of plasma membrane to lysosomes. Autophagy of normal (173) or hyperplastic smooth endoplasmic reticulum will supply the lysosomes with excessive lipids characteristic of this organelle. Proliferation of smooth endoplasmic reticulum (3,26,62,120,174) associated with enhanced drug metabolism of some AP drugs (112,174) has been observed. Other drugs, such as tilorone and DEAEH, are, however, known to decrease drug metabolism and phospholipid content of the microsomal fraction (103,175).

Structural Types of Myelinoid Bodies

Myelinoid bodies are secondary lysosomes arising from the processes of autophagy and heterophagy. Sequestered and phagocytosed proteins and carbohydrates are degraded, while certain lipids resist digestion, because they formed complexes with drugs either within the lysosomes or prior to their internalization into the lysosome. The drug-lipid complexes inside the lysosome are rearranged into lamellar, reticular or crystalloid structures to form concentric lamellar, reticular and crystalloid myelinoid bodies (26) (Figs. 6 and 7). Fusion of unicentric lamellar myelinoid bodies gives origin to multicentric myelinoid bodies. Myelinoid bodies containing parallel lamellae, also called "zebra bodies," are rarely induced by drugs but are common in lysosomal storage diseases (176). Concentric lamellae, reticular patterns and amorphous dense material may be present in the complex myelinoid body. The crystalloid myelinoid bodies contain phospholipids in regular densely spaced patterns, which are different (square, rectangular, linear, etc.) according to the plane of section (Fig. 7). They represent PLs in hexagonal phase (177). Their formation is favored by a high lysosomal content of phosphatidylethanolamine and of highly unsaturated fatty acids (126,177,178). Studies using *in vitro* models showed that the structural patterns of a

phospholipid depend on the chemical nature of the PL, the presence of fusogenic lipids, the temperature and hydration (179,180). Increased hydration is associated with a loose packing of lamellae.

Myelinoid bodies induced by chloroquine contain abundant clear matrix in addition to lamellae. The formation of such vacuolated myelinoid bodies has been attributed to osmotic swelling due to a rapid intralysosomal accumulation of drugs which are not easily protonized and therefore not absorbed to endoplasmic reticulum (41).

Myelinoid Bodies Unrelated to Cationic Amphiphilic Drugs

The epithelial cells of renal proximal tubules are known to accumulate heterogeneous dense bodies rich in myelin figures in response to a variety of stimuli which lead to an increased intake of PLs. Thus, e.g., an excessive internalization of plasma membrane is induced by resorption of proteins from the glomerular filtrate of old rats (181). Membranes of hemolyzed erythrocytes are incorporated by heterophagy (42). The strongly cationic aminoglycosides gentamycin and netilmicin bind to the brush border and are reabsorbed by endocytosis, accumulate in lysosomes, induce myelinoid bodies and accumulation of phospholipids (182-184). Paraquat is cationic in its dissolved form and may be similarly bound to acidic phospholipids to form renal DIM bodies (185). Lysosomes resembling myelinoid bodies are found in the cells of the renal papilla and inner medulla of potassium depleted rats and man (186) as a consequence of an inadequate intralysosomal metabolism of PLs.

Pathology

The accumulation of pulmonary foam cells and of myelinoid bodies may be associated with serious alterations of cellular functions and lead to clinical diseases. The pathological effect of AP drugs will depend on the type of cellular membranes to which the drugs are attached. Thus the formation of drug-lipid complexes in the plasma membrane may be expected to alter the membrane flow (187), inhibit membrane internalization (188) and phagocytosis (189). Chlorpromazine, triparanol and gentamycin inhibit ($\text{Na}^+ + \text{K}^+$) ATPase activity (190-192). AP drugs alter permeability of membranes (193,194). The properties of membranes are changed by an increase in the fluidity of the lipid (190,195-197). AP drugs cause transition of lipid acyl chain from organized gel to randomized liquid crystalline phase. The phase transition is associated with a lateral expansion, decrease in thickness and density of a membrane, and with altered function of membrane-bound proteins.

Attachment of the drug to endoplasmic reticulum may result in the most important functional changes (28). The alteration of activities of enzymes associated with membranes were discussed earlier. Presence of the

un-ionized drug in the membrane may be responsible for stabilization of lysosomes and of erythrocytes (157,198). Additional damage to membranes will depend on the metabolism of AP drugs into either stable epoxides or less stable arene oxides (199). Membranes containing drug-lipid complexes are recognized by the cell as abnormal, and are sequestered by the process of autophagy leading to the formation of DIM bodies.

Small numbers of myelinoid bodies are well tolerated by the cell. Large numbers may interfere with functional associations between organelles and impair metabolic processes. Cellular death may result for several reasons. The cell may be compressed in a limited space. The available pools of phospholipids needed for cell function and survival may be exhausted (200,201). Finally the cell may rupture into the lumen of a hollow structure, as seen in renal tubules (Fig. 8). Presence of myelin figures in the absence of cellular organelles, however, does not indicate cell rupture, but it represents the contents of myelinoid bodies extruded from the cell (202).

Functional impairment of overloaded macrophages may lead to a decreased resistance to infections (25). On the other hand, administration of amphiphilic drugs may be beneficial in controlling infections. Isolated alveolar macrophages from animals treated with AP drugs show an enhanced phagocytic activity and are more effective in killing bacteria (23). Chloroquine inhibits infections by enveloped viruses by preventing their uncoating within lysosomes (203). Chlorpromazine and trifluoperazine kill even phagocytized protozoa (204).

Amphiphilic drugs are known to induce adverse side effects in man. The foam cell syndrome was known to occur in Japanese patients since 1965. Yamamoto et al. (156) established that the syndrome is related to the intake of diethylaminoethoxyhexestrol and reproduced it in rats by the same drug. The term "drug-induced phospholipidosis" was introduced by Shikata et al. (205) and was popularized by Lüllmann-Rauch (41). It indicates a generalized lysosomal storage of phospholipids. The drug-induced lysosomal storage differs from inborn lysosomal storage diseases (27,67,76,101). It has not been established whether administration of AP drugs to patients with lysosomal storage diseases will have adverse effects because of an additional increase in lysosomal volume. A patient with a subclinical enzyme deficiency may be expected to develop a clinically significant disease.

Other clinical adverse effects induced by AP drugs in man are summarized in Table 4. Some of the serious adverse effects were discovered only after the drugs were used by man. The anorectic drugs have been associated with pulmonary hypertension in patients with familial predisposition (206). These vascular effects are related to an altered metabolism of serotonin rather than to foam cell accumulation (207). Chlorpheniramine was shown to decrease pulmonary clearance of 5-hydroxytryptamine (208). Most probably other AP drugs will have clinical side effects. It is therefore

imperative to search for the ability of each newly developed drug to induce myelinoid bodies in specific locations when given in large doses.

In animal experiments, massive accumulation of pulmonary foam cells is an excellent indicator of a drug-induced lysosomal storage. It should be remembered, however, that foam cells occur in other pathological entities (1) and in old normal rats (209). Preclinical screening of drugs in intact animals may not identify a drug as a myelinoid body inducer, if the drug is rapidly metabolized to an inactive form. This drug may, however, have adverse effects in a few patients who are unable to metabolize it. Such a drug may be recognized in *in vitro* systems. In performing *in vitro* studies, we should be aware of the fact that gentamycin used in cell cultures has been shown to be an inducer of DIM bodies (210). The work of Unsicker et al. (211) with lizards suggests, that screening methods utilizing cold-blooded animals are feasible.

Unique pharmacological properties of some AP drugs will necessitate their clinical use in spite of induction of DIM bodies. The adverse effects of such drugs will be more predictable and the clinical use will be safer provided that the principles of myelinoid body formation and degradation will be established in animal experiments. At first we need studies at cellular level on the interactions between AP drugs as well as studies on interactions of AP drugs with other drugs, food components and environmental contaminants. Only a few structural studies were reported on AP drug interactions (212-214). The type of dietary lipids in the food may affect the resorption, distribution, retention and metabolism of AP drugs. Thus, for example, cholesterol has been shown to antagonize the phospholipid-drug interaction *in vitro* (152).

Equally needed are studies on factors affecting the disappearance of myelinoid bodies after the discontinuation of treatment. DIM bodies in animal experiments and *in vitro* systems usually disappear in 1 to 10 weeks (33,62,178,212,215). The recovery in monkeys (216) and in man (33,72) may take years to be complete. While it is known that simultaneous administration of phenobarbital and chlorphentermine reduces the formation of alveolar foam cells and DIM bodies (132,146) experimental studies on factors enhancing the resolution of already accumulated DIM bodies and inclusions in alveolar foam cells are lacking.

Pulmonary histiocytosis induced by AP drugs deserves special consideration of additional factors, such as the amount of the surfactant available for phagocytosis, the metabolism of drugs by Clara cells (217) and the localization of drugs in the lung. The accumulation of foam cells could be enhanced by simultaneous administration of drugs and hormones (*L*-thyroxin) which increase the amounts of the surfactant within alveoli either by stimulating the secretion (218) or the proliferation of Type II pneumocytes (219). In this regard it is of interest that AP drugs are more toxic in hyperthyroid mice (220).

Drug interactions could also result from the mutual inhibition of amine accumulation in the lung, and reversible storage of drugs in the lung could precipitate toxic effects (133). Pathological investigations are needed to establish whether the adverse effects of AP drugs on the lungs are potentiated by simultaneous treatment with drugs known to localize and/or accumulate in the lung, such as amphetamine, chloramphetamine, desipramine, diphenhydramine, fluphenazine, methadone, propranolol and tripeleamine (221), and with drugs which inhibit the binding of chlorphen-

Table 4. Clinical adverse reactions associated with the administration of known inducers of myelinoid bodies.^a

Drug	Reactions ^a											References	
	Keratopathy	Retinopathy	Cataract	Neuropathy	Myopathy	Hepatomegaly	Splenomegaly	Liver fibrosis or cirrhosis	Hyperlipidemia	Pulmonary foam cells	Nephropathy		Malabsorption
Amiodarone	+			+						+			(41,121,252,296,305,306)
Chloroquine	+	+			+								(41,121)
Chlorpromazine	+		+										(121,297)
Diethylamino-ethoxyhexestrol						+	+	+	+	+			(33,95,156,205,242)
Gentamycin											+		(184,298)
Perhexiline				+		+		+					(124,125,275,307)
Quinacrine	+												(259)
Tamoxifen	+	+											(299,303,308)
Tilorone	+	+											(261,262)
Triparanol			+									+	(280,287,300,301)

^a + denotes reported.

termine to the lung, such as pronethalol, pyrilamine, *d*-propoxyphene and quinidine (134).

The chemicals which should be studied for their ability to modify the adverse effects of AP drugs and for their potential to induce myelinoid bodies under special circumstances include: the basic dyes, such as acridine orange or methylene blue; the lipophilic basic amines (221); and drugs with amphiphilic molecules, which were not fully studied. Priority should be given to drugs currently on the market such as the expectorant bromhexine; the antihistaminics carbinoxamine, cyproheptadine, diphenhydramine, diphenylpyraline, methapyrilene, pyrilamine and tripelenamine; the anticholinergics chlorphenoxamine and clofenetamine; the antidepressants butriptyline, desipramine, doxepine, protriptyline, trimipramine and zimelidine; the tranquilizers fluphenazine, promazine and triflupromazine; the analgesics methadone and *d*-propoxyphene; the gonad stimulating agents clomiphene and zuclophene; the suppressant of lupus erythematosus hydroxychloroquine; the anorectics phenpentermine and norfenfluramine; and the adrenergic blockers pronethacol and propranolol.

In clinical trials the peripheral blood is easily accessible for light microscopic and ultrastructural studies. Lymphocytes and plasma cells of animals and man respond to AP drugs by formation of vacuoles which correspond to myelinoid bodies (231,32,222). Some AP drugs such as perhexiline are, however, poor inducers of myelinoid bodies in lymphocytes (123).

The intra-alveolar accumulation of foamy macrophages is a form of pulmonary histiocytosis (7,20). Experimentally, alveolar foam cells can be induced by inhalation of aerosols of nickel oxide (223). Large PAS-positive cells can be induced by Freund's adjuvant (224) and are seen in tuberculous pneumonia of man (Fig. 9). These cells have a granular cytoplasm. Cells seen in lipid pneumonia, on the other hand, contain vacuoles of different sizes (Fig. 10). Massive accumulation of intra-alveolar histiocytes and hyperplasia of granular pneumocytes are found also in human and experimental desquamative interstitial pneumonia (225-230). These macrophages are, however, smaller, lack foamy appearance, and contain PAS-positive granules.

The drug-induced pulmonary histiocytosis (20) and desquamative pneumonia in man (231) may progress to alveolar proteinosis. The alveoli in alveolar proteinosis are filled with amorphous, proteinaceous, PAS-positive material which is rich in lipids and contains phospholipids (232-234). For this reason, the term alveolar lipoproteinosis is more exact (232). The intra-alveolar material seen in proteinoses of different origin is mostly derived from disintegrated macrophages (235-237) and from pneumocytes (232,233,238). Although the alveolar proteinosis induced by AP drugs in animals did not result in fibrosing alveolitis (21), cholesterol granulomata were observed in lungs of dogs treated with AY-9944 (1). Alveolar proteinosis is also observed in




FIGURE 9. Alveolar macrophages from a patient with tuberculous pneumonia. $\times 590$.




FIGURE 10. Alveolar macrophage from a patient with lipid pneumonia. $\times 590$.

acute silicosis (238,239) and as a response to a variety of nonspecific chemical irritants (235).

Diagnosis and Treatment

The alveolar proteinosis in man can be diagnosed in the electron microscope by examination of the sputum and of lung washings (240). Controlled volume bronchial lavage gives a good clinical response in treatment of alveolar proteinosis (240). Such treatment could be applied to clinically diagnosed cases of drug-induced alveolar histiocytosis. The degree of hepatic involvement in generalized lysosomal storage and the recovery can be followed by the levels of serum transaminases (72,124,212,241,242). The treatment of symptomatic generalized drug-induced lysosomal storage is discontinuation of the drug.

Terminology

Blohm (157) stressed the need for a standard nomenclature to facilitate indexing of drug-induced lipidoses.

We propose the following terms: drug-induced myelinoid body, DIM body: (a) concentric lamellar, (b) parallel lamellar, (c) reticular, (d) crystalloid, (e) mixed type; drug-induced foam cell; drug-induced lysosomal storage (lipidosis): (a) generalized, (b) organ-specific, e.g., renal, retinal, cerebrospinal (formerly cerebrospinal lipodystrophy, neuronal storage dystrophy), pulmonary (intra-alveolar histiocytosis).

Summary

Administration of amphiphilic drugs to experimental animals causes formation of myelinoid bodies in many cell types, accumulation of foamy macrophages in pulmonary alveoli, and pulmonary alveolar proteinosis. These changes are the result of an interaction between the drugs and phospholipids which leads to an alteration in physicochemical properties of the phospholipids. Impairment of the digestion of altered pulmonary secretions in phagosomes of macrophages results in accumulation of foam cells in pulmonary alveoli. Impairment of the metabolism of altered phospholipids removed by autophagy induces an accumulation of myelinoid bodies. The administration of amphiphilic compounds thus causes pulmonary intra-alveolar histiocytosis which is a part of a drug-induced lysosomal storage or generalized lipidosis.

NOTE ADDED IN PROOF: Amphiphilic drugs have been the subject of numerous recent studies not otherwise cited in this review (309–340).

Figure 2 is reproduced with permission of the Academic Press. The secretarial assistance of Mrs. Betty Jane Belisle and Mrs. Barbara Ro is gratefully acknowledged.

REFERENCES

- Greselin, E. An inhibitor of cholesterol biosynthesis and the alveolar macrophages. *Can J. Comp. Med. Vet. Sci.* 30: 121–126 (1966).
- Lüllmann-Rauch, R., and Reil, G. H. Fenfluramine-induced ultrastructural alterations in tissues of rats and guinea pigs. *Naunyn Schmiedeberg's Arch. Pharmacol.* 285: 175–184 (1974).
- Lüllmann-Rauch, R., and Scheid, D. Intraalveolar foam cells associated with lipidosis-like alterations in lung and liver of rats treated with tricyclic psychotropic drugs. *Virchows Arch. (Cell Pathol.)* 19: 255–268 (1975).
- Parwaresch, M. R., Reil, G. H., and Seiler, K. U. Studies on animal and organ specificity of chlorphentermine induced morphological alterations. *Res. Exp. Med. (Berlin)* 161: 272–288 (1973).
- Magnusson, G., and Magnusson, O. Clofex-induced pulmonary changes in rats. *Beitr. Pathol.* 146: 79–88 (1972).
- Hruban, Z., and Slesers, A. Alterations in outer renal medulla produced by chloroquine. *Fed. Proc.* 30: 637 (1971).
- Hruban, Z., Slesers, A., and Aschenbrenner, I. Pulmonary intraalveolar histiocytosis induced by drugs. *Toxicol. Appl. Pharmacol.* 26: 72–85 (1973).
- Kikkawa, Y., and Suzuki, K. Alterations of cellular and acellular alveolar and bronchiolar walls produced by hypocholesteremic drug AY 9944. *Lab. Invest.* 26: 441–447 (1972).
- Lüllmann, H., Lüllmann-Rauch, R., and Reil, G. H. A comparative ultrastructural study of the effects of chlorphentermine and triparanol in rat lung and adrenal gland. *Virchows Arch. (Cell Pathol.)* 12: 91–103 (1973).
- Lüllmann-Rauch, R., Reil, G. H., Rossen, E., and Seiler, K. U. The ultrastructure of rat lung changes induced by an anorectic drug (chlorphentermine). *Virchows Arch. (Cell Pathol.)* 11: 167–181 (1972).
- Karabelnik, D., Zbinden, G., and Baumgartner, E. Drug-induced foam cell reactions in rats. I. Histopathologic and cytochemical observations after treatment with chlorphentermine, RMI 10.393 and RO 4-4318. *Toxicol. Appl. Pharmacol.* 27: 395–407 (1974).
- Staiger, G. R. Light-microscopic demonstration of drug-induced myelin bodies in the liver of rats. *Experientia* 30: 385–386 (1974).
- Gräbner, R., and Meerbach, W. Morphological and biochemical alterations of the lung after application of chlorphentermine. *Exptl. Pathol.* 17:303–311 (1979).
- Vijayaratnam, G. S., and Corrin, B. Fine structural alterations in the lungs of iprindole-treated rats. *J. Pathol.* 114: 233–239 (1974).
- Reasor, M. J., Massey, C. A., Koshut, R. A., and Castranova, V. Multinucleation in alveolar macrophages from rats treated with chlorphentermine. *Lab. Invest.* 46: 224–230 (1982).
- Vijayaratnam, G. S., and Corrin, B. Origin of the pulmonary alveolar macrophage studied in the iprindole treated rat. *J. Pathol.* 108: 115–113 (1972).
- Flodh, H., and Magnusson, G. Genesis of foam cells: study in rats after administration of clofex. *Virchows Arch. (Cell Pathol.)* 12: 360–366 (1973).
- Corrin, B., and King, E. Experimental endogenous lipid pneumonia and silicosis. *J. Pathol.* 97: 325–330 (1969).
- Smith, P., Heath, D., and Hasleton, P. S. Electron microscopy of chlorphentermine lung. *Thorax* 28: 559–566 (1973).
- Vijayaratnam, G. S., and Corrin, B. Pulmonary alveolar proteinosis developing from desquamative interstitial pneumonia in long term toxicity studies in the rat. *Virchows Arch. (Pathol. Anat.)* 358: 1–10 (1973).
- Smith, P., Heath, D., and Hasleton, P. Effects of prolonged administration of chlorphentermine on the rat lung. *Pathol. Europ.* 9: 273–287 (1974).
- Bockhardt, H., and Lüllmann-Rauch, R. Zimelidine-induced lipidosis in rats. *Acta Pharmacol. Toxicol.* 47: 45–48 (1980).
- Reasor, M. J. Drug-induced lipidosis and the alveolar macrophage. *Toxicology* 20: 1–33 (1981).
- Wold, J. S., Joost, R. R., Griffing, W. J., Marroquine, P., and Harris, P. N. Phospholipid accumulation in rats produced by fluoxetine and chlorphentermine. *Toxicol. Appl. Pharmacol.* 37: 118–119 (1976).
- Hruban, Z., Swift, H., and Slesers, A. Effect of triparanol and diethanolamine on the fine structure of hepatocytes and pancreatic acinar cells. *Lab. Invest.* 14: 1652–1672 (1965).
- Hruban, Z., Slesers, A., and Hopkins, E. Drug-induced and naturally occurring myeloid bodies. *Lab. Invest.* 27: 62–70 (1972).
- Lüllmann, H., Lüllmann-Rauch, R., and Wassermann, O. Drug-induced phospholipidoses. *CRC Crit. Rev. Toxicol.* 4: 185–218 (1975).
- De La Iglesia, F. A., Feuer, G., McGuire, E. J., and Takada, A. Morphological and biochemical changes in the liver of various species in experimental phospholipidosis after diethylaminoethoxyhexestrol treatment. *Toxicol. Appl. Pharmacol.* 34: 28–44 (1975).
- Ghadially, F. N. *Ultrastructural Pathology of the Cell*. Butterworths, London and Boston, 1975.
- Gray, J. E., Weaver, R. N., Stern, K. F., and Phillips, W. A. Foam cell response in the lung and lymphatic tissues during long-term high-level treatment with erythromycin. *Toxicol. Appl. Pharmacol.* 45: 701–711 (1978).
- Imoto, S., Hiraide, S., Mitsutani, S., Matsumoto, H., Kaneko, S., Miyake, T., Akasaka, K., Shinji, Y., Yamamoto, A., Adachi, S., Fukui, I., Sasaki, H., Mori, Y., and Okada, S. A case of

- drug-induced lipidosis with "sea-blue histiocytes" in the bone marrow. Japan. Arch. Int. Med. 18: 83-108 (1971).
32. Kitani, T., Yamamoto, A., Adachi, S., Tako, H., Imanaka, T., Nasut, T., and Nishikawa, M. Drug-induced lipidosis (IV). Morphological aspect of leucocytes and bone marrow cells in human cases and animal experiments induced by 4,4'-diethylaminoethoxyhexestrol. Acta Haematol. Japan 35: 131-145 (1972).
 33. Akeda, S. A study on the lipidosis induced by a coronary vasodilator, 4,4'-diethylaminoethoxyhexestrol dihydrochloride. Mie Medical J. 22: 65-96 (1972).
 34. Gautier, M., Raulin, J., LaPous, D., Lorient, C., Carreau, J. P., and Scoto, J. Children sea-blue histiocytosis (2 cases) compared with phospholipidosis induced by 4-4'DET (4,4'-p(diethylamino-2-ethoxyphenyl)3,4-hexane) in rat. Biomedicine 26: 52-60 (1977).
 35. Hruban, Z., Spargo, B., Swift, H., Wissler, R. W., and Kleinfeld, R. Focal cytoplasmic degradation. Am. J. Pathol. 42: 657-683 (1963).
 36. Goren, M. B., Hart, P. D., Young, M. R., and Armstrong, J. A. Prevention of phagosome-lysosome fusion in cultured macrophages by sulfatides of mycobacterium tuberculosis. (Proc. Natl. Acad. Sci. (U.S.)) 73: 2510-2514 (1976).
 37. Hart, P. D. A. Phagosome-lysosome fusion in macrophages: a hinge in the intracellular fate of ingested microorganisms. In: Lysosomes in Biology and Pathology (J. T. Dingle, P. J. Jacques and I. H. Shaw, Eds.), Vol. 6, North-Holland, New York, 1979, pp. 409-423.
 38. Mason, R. J., and Williams, M. C. Phospholipid composition and ultrastructure of A 549 cells and other cultured pulmonary epithelial cells of presumed type II cell origin. Biochim. Biophys. Acta 617: 36-50 (1980).
 39. Drenckhahn, D. Anterior polar cataract and lysosomal alterations in the lens of rats treated with the amphiphilic lipidosis-inducing drugs chloroquine and chlorpheniramine. Virchows Arch. (Cell Pathol.) 27: 255-266 (1978).
 40. Jung, H. J., and Suzuki, K. Morphological changes in CNS of rats treated with perhexiline maleate (Pexid). Acta Neuropathol. (Berlin) 42: 159-164 (1978).
 41. Lüllmann-Rauch, R. Drug-induced lysosomal storage disorders. In: Lysosomes in Biology and Pathology (J. T. Dingle, P. J. Jacques and I. H. Shaw, Eds.), Vol. 6, North-Holland, New York, 1979, pp. 49-130.
 42. Torhorst, J., Richter, L., and Rohr, H. P. Entwicklung und Umwandlung lysosomaler Funktionsformen unter besonderer Berücksichtigung des glatten endoplasmatischen Reticulum. Virchows Arch. 343: 64-74 (1967).
 43. Quennedy, A., and Brossut, R. Les glandes mandibulaires de *Blaberus craniifer* burm. (Dictyoptera, Blaberidae) development, structure et fonctionnement. Tissue Cell 7: 503-517 (1975).
 44. Riley, J., James, J. L., and Banaja, A. A. The possible role of the frontal and sub-parietal gland systems of the pentastomid *Reighardia sterna* (Diesing, 1864) in the evasion of the host immune response. Parasitology 78: 53-66 (1979).
 45. Henkart, M. Light induced changes in the structure of pigmented granules in Aplysia neurons. Science 188: 155-157 (1975).
 46. Sohal, R. S., Peters, P. D., and Hall, T. A. Origin, structure, composition and age-dependence of mineralized dense bodies (concretions) in the mid-gut epithelium of the adult housefly *Musca domestica*. Tissue Cell 9: 87-102 (1977).
 47. Kühnel, W. The glandulae rectales of the rabbit. Z. Zellforsch. 122: 574-583 (1971).
 48. Brunck, V., Ericsson, J. L. E., Ponton, J., and Westermarck, B. Residual bodies and "aging" in cultured human glia cells. Exptl. Cell Res. 79: 1-14 (1973).
 49. Vaughan, D. W. Membranous bodies in the cerebral cortex of aging rats: An electron microscope study. J. Neuropathol. Exptl. Neurol. 35: 152-166 (1976).
 50. Del Cerro, M. P., and Snyder, R. S. Studies on dilantin intoxication. Ultrastructural analogies with the lipidosis. Neurology 17: 452-466 (1967).
 51. Shinozuka, H., Reid, I. M., Shull, K. H., Liang, H., and Farber, E. Dynamics of liver cell injury and repair. I. Lab. Invest. 23: 253-267 (1970).
 52. Suzuki, K., and De Paul, L. D. Cellular degeneration in developing central nervous-system of rats produced by hypocholesterolemic drug AY 9944. Lab. Invest. 25: 546-555 (1971).
 53. Le Beux, Y., Hetenyi, G., and Philips, M. J. Mitochondrial myelin-like figures: a non-specific reactive process of mitochondrial phospholipid membranes to several stimuli. Z. Zellforsch. 99: 491-506 (1969).
 54. Blanchette-Mackie, E. J., and Scow, R. O. Retention of lipolytic products in chylomicrons incubated with lipoprotein lipase: electron microscope study. J. Lipid Res. 17: 57-67 (1976).
 55. Blanchette-Mackie, E. J., and Ucci, A. A. Lamellar structure and lipolytic activity in hepatic intracellular lipid inclusions. J. Cell Biol. 70: 285a (1976).
 56. Sun, C. N., White, H. J., and Lucas, G. Crystalloid inclusions in the Schwann cell cytoplasm. Beitr. Pathol. 148: 402-406 (1973).
 57. Stock, C., and Grenier, J. F. Intracellular and extracellular myelin bodies in exocrine pancreas of obese hyperglycaemic mouse. Cytobiologie 12: 111-118 (1975).
 58. Abraham, R., and Hendy, R. Effects of chronic chloroquine treatment on lysosomes of rat liver cells. Exptl. Mol. Pathol. 12: 148-159 (1970).
 59. Zwahlen, R., Richardson, B. P., and Hauser, R. E. The production and elimination of myeloid bodies by cultured pancreatic islet cells. J. Ultrastruct. Res. 67: 340-356 (1979).
 60. Hildebrand, J., Thys, O., and Gérin, Y. Alterations of rat liver lysosomes and smooth endoplasmic reticulum induced by the diazafluoranthene derivative AC-3579. II. Effects of the drug on phospholipid metabolism. Lab. Invest. 28: 83-86 (1979).
 61. Hoenig, V., and Werner, F. Effect of perhexiline maleate on lipid metabolism in the rat. Arzneim. Forsch. 29: 1395-1398 (1979).
 62. Thys, O., Hildebrand, J., Gérin, Y., and Jacques, P. J. Alterations of rat liver lysosomes and smooth endoplasmic reticulum induced by the diazafluoranthene derivative AC-3579. I. Morphologic and biochemical lesions. Lab. Invest. 28: 70-82 (1973).
 63. Plantavid, M., Chap, H., Lloveras, J., and Douset-Blazy, L. Cationic amphiphilic drugs as a potential tool for modifying phospholipids of tumor cells. An in vitro study of chlorpromazine effects on Krebs II ascites cells. Biochem. Pharmacol. 30: 293-297 (1981).
 64. Matsuzawa, Y., and Hostetler, K. Y. Effects of chloroquine and 4,4'-bis(diethylaminoethoxy)- α,β -diethyldiphenylethane on the incorporation of [3 H]glycerol into the phospholipids of rat liver lysosomes and other subcellular fractions, in vivo. Biochim. Biophys. Acta 620: 592-602 (1980).
 65. Lageron, A., and Maziere, J. C. Histochemical and biochemical effects of homochlorocyclizine on liver. J. Clin. Chem. Clin. Biochem. 19: 745 (1981).
 66. Blok, J., Mulder-Stapel, A. A., Ginsel, L. A., and Daems, W. T. The effect of chloroquine on lysosomal function and cell-coat glycoprotein transport in the absorptive cells of cultured human small-intestine tissue. Cell Tissue Res. 218: 227-251 (1981).
 67. Klinghardt, G. W., Fredman, P., and Svennerholm, L. Chloroquine intoxication induces ganglioside storage in nervous tissue: A chemical and histopathological study of brain, spinal cord, dorsal root ganglia, and retina in the miniature pig. J. Neurochem. 37: 897-908 (1981).
 68. Nilsson, O., Fredman, P., Klinghardt, G. W., Dreyfus, H., and Svennerholm, L. Chloroquine-induced accumulation of gangliosides and phospholipids in skeletal muscles. Eur. J. Biochem. 116: 565-571 (1981).
 69. Fredman, P., Klinghardt, G. W., Nilsson, O., and Svennerholm, L. Lipid accumulation in liver, spleen, lungs and kidneys of miniature-pigs after chloroquine treatment. Biochem. J. 201: 581-588 (1982).
 70. Adachi, M., Tsai, C. Y., Greenbaum, M., Mask, B., and Volk, B. W. Ultrastructure and biochemical studies of rat CNS and viscera after subcutaneous injection of chlorpheniramine. In: Current Trends in Sphingolipidoses and Allied Disorders (B. W. Volk and L. Schnek, Eds.), Plenum Press, New York, 1976, pp. 429-451.
 71. Pappu, A. S., and Hauser, G. Alterations of phospholipid

- metabolism in rat cerebral cortex mince induced by cationic amphiphilic drugs. *J. Neurochem.* 37: 1006-1014 (1981).
72. Lageron, A., Poupon, R., De Saint-Maur, P. P., and Levy, V. G. Liver ganglioside storage after perhexiline maleate. *Lancet* i: 483 (1977).
 73. Hauw, J. J., Mussini, J. M., Boutry, J. M., Escourolle, R., Pollet, S., Albouz, S., Harpin, M. L., and Baumann, N. Perhexiline maleate induced lipidosis in human peripheral nerve and tissue culture: ultrastructural and biochemical changes. *Clin. Toxicol.* 18: 1405-1409 (1981).
 74. Hauw, J. J., Boutry, J. M., Albouz, S., Harpin, M. L., Baudrimont, M., Escourolle, R., and Baumann, N. Perhexiline maleate-induced lipidosis in cultured human fibroblasts: cell kinetics, ultrastructural and biochemical studies. *Virchows Arch. (Cell Pathol.)* 34: 239-249 (1980).
 75. Hill, P., and Greselin, F. Lung phospholipids: biochemical and histologic changes induced by the cholesterol biosynthesis inhibitor AY-9944. *Toxicol. Appl. Pharmacol.* 11: 245-256 (1967).
 76. Matsuzawa, Y., and Hostetler, K. Y. Studies on drug-induced lipidosis: subcellular localization of phospholipid and cholesterol in the liver of rats treated with chloroquine or 4,4'-bis(diethylaminoethoxy)- α,β -diethylidiphenylethane. *J. Lipid Res.* 21: 202-214 (1980).
 77. Dietert, S. E., and Scallen, T. J. An ultrastructural and biochemical study of the effects of three inhibitors of cholesterol biosynthesis upon murine adrenal gland and testis. *J. Cell Biol.* 40: 44-60 (1969).
 78. Yates, R. D., Arai, K., and Rappoport, D. A. Fine structure and chemical composition of opaque cytoplasmic bodies of triparanol-treated Syrian hamsters. *Exptl. Cell Res.* 47: 459-478 (1967).
 79. May, C., Doriaux, M., Thys, O., Hildebrand, J., and Drochmans, P. Effects of the diazafluoranthene derivative AC-3579 on rat liver. A. Isolation and characterization of lamellate cytosomes and subcellular localization of the drug. *Biol. Cellulaire* 34: 187-196 (1979).
 80. Chatelain, P., and May, C. Abnormal microviscosity of lamellate cytosomes induced by a diazafluoranthene derivative. *Biochem. Pharmacol.* 28: 3467-3470 (1979).
 81. Seiler, K. U., and Wassermann, O. Drug induced phospholipidosis. II. Alterations in the phospholipid pattern of organs from mice, rats and guinea pigs after chronic treatment with chlorphentermine. *Naunyn Schmiedeberg's Arch. Pharmacol.* 288: 261-268 (1975).
 82. Gloster, J., Heath, D., Haselton, P., and Harris, P. Effect of chlorphentermine on the lipids of rat lungs. *Thorax* 31: 558-564 (1976).
 83. Karabelnik, D., Hörler, D. A., and Zbinden, G. Lipid distribution pattern in lungs and alveolar macrophages of rats treated with 3 foam cell-inducing drugs. *Proc. Europ. Soc. Toxicol.* 17: 315-321 (1975).
 84. Karabelnik, D., and Zbinden, G. Drug-induced foam cell reactions in rats. II. Chemical analysis of lipids stored in lungs and foam cells after treatment with chlorphentermine, 5-[p-(fluoren-9-ylidenemethyl)phenyl]-2-piperidineethanol (RMI 10.393) and 1-chloramitriptyline. *Hoppe Seyler's Z. Physiol. Chem.* 356: 1151-1160 (1975).
 85. Brody, A. R., Clay, M. F., Collins, M. M., Eden, A. C., and McDermott, M. Effects of chlorphentermine hydrochloride on the surface tension properties of the rat lung. *J. Physiol. (London)* 245: 105P-106P (1975).
 86. Hruban, Z., Rubenstein, A. H., and Slesers, A. Alterations in pancreatic beta cells induced by cyclizine. *Lab. Invest.* 26: 270-277 (1972).
 87. Frisch, W., and Lüllmann-Rauch, R. Effects of several lipidosis-inducing drugs upon the area postrema and adjacent medullary nuclei of adult rats. *Acta Neuropathol.* 52: 179-187 (1980).
 88. Reasor, M. J., Koshut, R. A., and Castranova, V. Biochemical characteristics of rat alveolar macrophages with chlorphentermine-induced phospholipidosis: variations with increasing cell size. *Exptl. Mol. Pathol.* 31: 297-307 (1979).
 89. Schmien, R., Seiler, K. U., and Wassermann, O. Drug-induced phospholipidosis. I. Lipid composition and chlorphentermine content of rat lung tissue and alveolar macrophages after chronic treatment. *Naunyn Schmiedeberg's Arch. Pharmacol.* 283: 331-334 (1974).
 90. Reasor, M. J. Cumene hydroperoxide-mediated lipid peroxidation in rat alveolar macrophages following induction of phospholipidosis with chlorphentermine. *Toxicology* 18: 159-168 (1980).
 91. Defrise-Quertain, F., Chatelain, P., and Ruyschaert, J. M. Phospholipase inactivation induced by an aminopiperazine derivative: A study at the lipid-water interface. *J. Pharm. Pharmacol.* 30: 608-612 (1978).
 92. Spangler, W. L., Adelman, R. D., Conzelman, G. M., and Ishizaki, G. Gentamycin nephrotoxicity in the dog: Sequential light and electron microscopy. *Vet. Pathol.* 17: 206-217 (1980).
 93. Adachi, S., Matsuzawa, Y., Yokomura, T., Ishikawa, K., Uhara, S., Yamamoto, A., and Nishikawa, M. Studies on drug-induced lipidosis. (V). Changes in the lipid composition of rat liver and spleen following the administration of 4,4'-dimethylaminoethoxyhexestrol. *Lipids* 7: 1-7 (1972).
 94. Tjong, H. B., and Debuch, H. Lysosomal bis(monoacylglycerol)-phosphate of rat liver, its induction by chloroquine and its structure. *Hoppe-Seyler's Z. Physiol. Chem.* 359: 71-79 (1978).
 95. Yamamoto, A., Adachi, S., Ishibe, T., Shinji, Y., Kaki-uchi, Y., Seki, K., and Kitani, T. Accumulation of acidic phospholipids in a case of hyperlipidemia with hepatosplenomegaly. *Lipids* 5: 566-571 (1970).
 96. Yamamoto, A., Adachi, S., Kitani, T., Shinji, Y., Seki, K., Nasu, T., and Nishikawa, M. Drug-induced lipidosis in human cases and in animal experiments. *J. Biochem.* 69: 613-615 (1971).
 97. Yamamoto, A., Adachi, S., Matsuzawa, Y., Kitani, T., Hiraoka, A., and Seki, K. Studies on drug-induced lipidosis. VII. Effects of bis- β -diethylaminoethylether of hexestrol, chloroquine, homochlorcyclizine, prenylamine, and diazcholesterol on the lipid composition of rat liver and kidney. *Lipids* 11: 616-622 (1976).
 98. Matsuzawa, Y., Yamamoto, A., Adachi, S., and Nishikawa, M. Studies on drug-induced lipidosis. VIII. Correlation between drug accumulation and acidic phospholipids. *J. Biochem.* 82: 1369-1377 (1977).
 99. Fedorko, M. Effect of chloroquine on morphology of cytoplasmic granules in maturing human leucocytes: an ultrastructural study. *J. Clin. Invest.* 46: 1932-1942 (1967).
 100. Kalina, M., and Bubis, J. J. Myeloid bodies formation in triparanol treated cultured cells. *Virchows Arch. (Cell Pathol.)* 19: 349-357 (1975).
 101. Stäubli, W., Schweizer, W., and Suter, J. Some properties of myeloid bodies induced in rat liver by an antidepressant drug (Maprotiline). *Exptl. Mol. Pathol.* 28: 177-195 (1978).
 102. Stäubli, W., Schweizer, W., Suter, J., and Hess, R. Ultrastructural and biochemical action of benzoctamine and maprotiline on the rat liver. *Agents Actions* 4: 391-403 (1974).
 103. De La Iglesia, I. A., Feuer, G., Takada, A., and Matsuda, Y. Morphologic studies on secondary phospholipidosis in human liver. *Lab. Invest.* 30: 539-549 (1974).
 104. Kasama, K., Yoshida, K., Takeda, S., Tsujimura, R., and Hasegawa, S. Inhibition of acid esterase in rat liver by 4,4'-diethylaminoethoxyhexestrol. *Lipids* 11: 718-721 (1976).
 105. Lüllmann, H., and Möisinger, E. U. Increased lysosomal enzyme activities in tissues of rats suffering from chlorphentermine induced lipidosis. *Biochem. Pharmacol.* 28: 1015-1016 (1979).
 106. Ramsey, R. B., and Fischer, V. W. The biochemical and morphological response of hydrolytic enzymes in the developing brain to hypocholesterolemic agents. *Acta Neuropathol. (Berlin)* 49: 89-94 (1980).
 107. Ridout, R. M., Decker, R. S., and Wildenthal, K. Chloroquine-induced lysosomal abnormalities in cultured foetal mouse hearts. *J. Mol. Cell Cardiol.* 10: 175-183 (1978).
 108. Watts, S. D. M., Orpin, A., and MacCormick, C. Lysosomes and tegument pathology in the chemotherapy of schistosomiasis with 1,7-bis(p-aminophenoxy)heptane (153C51). *Parasitology* 78: 287-294 (1979).
 109. Gaton, E., and Wolman, M. Histochemical study on the pathogenesis of chlorcyclizine-induced pulmonary lipidosis. *Histochemistry* 63: 203-207 (1979).

110. Reasor, M. J., Trush, M. A., and Walker, E. R. Changes in lysosomal properties of alveolar macrophages of rats treated with chlorphentermine. *Toxicol. Appl. Pharmacol.* 46: 261-264 (1978).
111. Barnhart, J. W. The effect of chlorcyclizine on cholesterol metabolism in mice. *Toxicol. Appl. Pharmacol.* 27: 449-455 (1974).
112. Burns, J. J. Drug metabolism and teratogenic studies. In: *Environmental Variables in Oral Disease* (S. J. Kreshover and F. J. McClure, Eds.), American Association for Advancement of Science, Washington, DC, 1966, pp. 259-271.
113. Corte, L. D., Gremigni, D., Megazzini, I., Mobilio, R., and Sgaragli, G. P. Chlorimipramine-induced phospholipidosis: Biochemical and pharmacokinetic observations. *Brit. J. Pharmacol.* 67: 449P-450P (1979).
114. Eichelbaum, M., Hengstmann, J. H., and Dengler, H. J. Das Verteilungsmuster des Chlorphentermins bei Ratte, Kaninchen und Schwein. *Naunyn-Schmiedeberg's Arch. Pharmacol. Exptl. Pathol.* 267: 446-456 (1970).
115. Kuntzman, R., Klutch, A., Tsai, I., and Burns, J. J. Physiological distribution and metabolic inactivation of chlorcyclizine and cyclizine. *J. Pharmacol. Exptl. Therap.* 149: 29-35 (1965).
116. Narrod, S. A., Wilk, A. L., and King, C. T. G. Metabolism of meclizine in the rat. *J. Pharmacol. Exptl. Therap.* 147: 380-384 (1965).
117. Matsuzawa, Y., Yokomura, T., Ishikawa, K., Adachi, S., and Yamamoto, A. Studies on drug-induced lipidosis VI. Identification and determination of the drug and its metabolite in lipidosis induced by 4,4'-diethylaminoethoxyhexestrol. *J. Biochem.* 72: 615-621 (1972).
118. Minchin, R. F., Ilett, K. F., and Madsen, B. W. Chlorphentermine binding in rat lung subcellular fractions and its displacement by desmethylimipramine. *Biochem. Pharmacol.* 28: 2273-2278 (1979).
119. Chatelain, P., Berliner, C., Ruysschaert, J. M., and Jaffe, J. Effect of a diazafluoranthene derivative on phospholipases. A study at the air-water interface. *Biochim. Biophys. Acta* 419: 540-546 (1976).
120. Laurent, G., Hildebrand, J., and Thys, O. Alterations of rat liver lysosomes and smooth endoplasmic reticulum induced by the diazafluoranthene derivative AC-3579. III. Mechanism and site of action. *Lab. Invest.* 32: 580-584 (1975).
121. Seiler, K. U., Thiel, H. J., and Wassermann, O. Die Chloroquinkerneratopathie als Beispiel einer arzneimittelinduzierten Phospholipidosis. *Klin. Monatsbl. Augenheilkd.* 170: 64-73 (1977).
122. Allan, D., and Michell, R. H. Enhanced synthesis de novo of phosphatidylinositol in lymphocytes treated with cationic amphiphilic drugs. *Biochem. J.* 148: 471-478 (1975).
123. Lüllmann, H., and Lüllmann-Rauch, R. Perhexiline induces generalized lipidosis in rats. *Klin. Wochenschr.* 56: 309-310 (1978).
124. Beaugrand, M., Poupon, R., Levy, V. G., Gillard, P., Lageron, A., Lecomte, D., Darnis, F., and Ferrier, J. P. Lesions hépatiques dues au maléate de perhexiline. Étude clinique, biologique, histopathologique, ultrastructurale et biochimique de 7 cas. *Gastroenterol. Clin. Biol.* 2: 579-588 (1978).
125. Pessayre, D., Bichara, M., Feldmann, G., Degott, C., Potet, F., and Benhamou, J. P. Perhexiline maleate-induced cirrhosis. *Gastroenterology* 76: 171-177 (1979).
126. Drenckhahn, D., and Lüllmann-Rauch, R. Drug-induced retinal lipidosis: differential susceptibilities of pigment epithelium and neuroretina toward several amphiphilic cationic drugs. *Exptl. Mol. Pathol.* 28: 360-371 (1978).
127. Lüllmann, H., and Wehling, M. The binding of drugs to different polar lipids *in vitro*. *Biochem. Pharmacol.* 28: 3409-3415 (1979).
128. Goormaghtigh, E., Chatelain, P., Caspers, J., and Ruysschaert, J. M. Evidence of a specific complex between adriamycin and negatively-charged phospholipids. *Biochim. Biophys. Acta* 597: 1-14 (1980).
129. Klinghardt, G. W. Neuronal storage dystrophy in chronic chloroquine intoxication. *Neurotoxicology* 1: 371-380 (1977).
130. Bickel, M. H., and Minder, R. Metabolism and biliary excretion of the lipophilic drug molecules, imipramine and desmethyl-imipramine in the rat. II. Uptake into bile micelles. *Biochem. Pharmacol.* 19: 2437-2443 (1970).
131. Hook, G. E. R., and Bend, J. R. Pulmonary metabolism of xenobiotics. *Life Sci.* 18: 279-290 (1976).
132. Kacev, S., Narbaitz, R., Ruddick, J. A., and Villeneuve, D. C. Role of drug metabolism in protection against chlorphentermine-induced pulmonary phospholipidosis in adult rats. *Exptl. Mol. Pathol.* 35: 98-107 (1981).
133. Anderson, M. W., Orton, T. C., Pickett, R. D., and Eling, T. E. Accumulation of amines in the isolated perfused rabbit lung. *J. Pharmacol. Exptl. Therap.* 189: 456-466 (1974).
134. Minchin, R. F., Madsen, B. W., and Ilett, K. F. Effect of desmethylimipramine on the kinetics of chlorphentermine accumulation in isolated perfused rat lung. *J. Pharmacol. Exptl. Therap.* 211: 514-518 (1979).
135. Wilson, A. G. E., Pickett, R. D., Eling, T. E., and Anderson, M. W. Studies on the persistence of basic amines in the rabbit lung. *Drug Metab. Dispos.* 7: 420-424 (1979).
136. Tjälve, H., Olsson, S., and Andersson, A. The uptake of ¹⁴C-chloroquine by mouse pancreatic islets *in vitro*. *Acta Pharmacol. Toxicol.* 47: 38-44 (1980).
137. Angevine, L. S., Ohmiya, Y., and Mehendale, H. M. Effect of chlorphentermine pretreatment on chlorphentermine uptake by isolated perfused rat lung. *Drug Metab. Dispos.* 10: 68-73 (1982).
138. Buckingham, S., Heinemann, H. O., Sommers, S. C., and McNary, W. I. Phospholipid synthesis in the large pulmonary alveolar cell. *Am. J. Pathol.* 48: 1027-1041 (1966).
139. Hallman, M., and Gluck, L. Phosphatidylglycerol in lung surfactant. II. Subcellular distribution and mechanism of biosynthesis *in vitro*. *Biochim. Biophys. Acta* 409: 172-191 (1975).
140. Naimark, A. Cellular dynamics and lipid metabolism in the lung. *Fed. Proc.* 32: 1967-1971 (1973).
141. Desai, R., Tetley, T. D., Curtis, C. G., Powell, G. M., and Richards, R. J. Studies on the fate of pulmonary surfactant in the lung. *Biochem. J.* 176: 455-462 (1978).
142. Karabelnik, D., and Zbinden, G. Palmitic acid-1-¹⁴C incorporation and turnover in lung phospholipids of rats treated with chlorphentermine, RMI 10.393 and RO 4-4318. *Arch. Toxicol.* 35: 163-174 (1976).
143. Lüllmann-Rauch, R. Chlorphentermine-induced ultrastructural alterations in foetal tissues. *Virchows Arch. (Cell Pathol.)* 12: 295-302 (1973).
144. Kikkawa, Y., and Motoyama, E. K. Effect of AY-9944, a cholesterol biosynthesis inhibitor on fetal lung development and on the development of type II alveolar epithelial cells. *Lab. Invest.* 28: 48-54 (1973).
145. Kacew, S., Narbaitz, R., and Dubas, T. C. Biochemical and morphologic investigation of the influence of chlorphentermine and subsequent withdrawal on newborn rat lung. *Toxicol. Appl. Pharmacol.* 47: 185-191 (1979).
146. Kacew, S., and Narbaitz, R. The effect of phenobarbital on chlorphentermine-induced lipidosis-like alterations in renal tissue of adult and newborn rats. *Virchows Arch. (Cell Pathol.)* 36: 59-63 (1981).
147. Brindley, D. N., Allan, D., and Michell, R. H. The redirection of glyceride and phospholipid synthesis by drugs including chlorpromazine, fenfluramine, imipramine, mepyrmine and local anaesthetics. *J. Pharm. Pharmacol.* 27: 462-464 (1975).
148. Lüllmann, H., Plösch, H., and Ziegler, A. Ca replacement by cationic amphiphilic drugs from lipid monolayers. *Biochem. Pharmacol.* 29: 2969-2974 (1980).
149. Michell, R. H., Allan, D., Bowley, M., and Brindley, D. N. A possible metabolic explanation for drug-induced phospholipidosis. *J. Pharm. Pharmacol.* 28: 331-332 (1976).
150. Schwarting, H., Seiler, K. U., and Wassermann, O. On the mechanism of drug-induced phospholipidosis. *Naunyn-Schmiedeberg's Arch. Pharmacol. (Suppl.)* 293: R57 (1976).
151. Seydel, J. K., and Wassermann, O. NMR-studies on the molecular basis of drug-induced phospholipidosis. *Naunyn-Schmiede-*

- bergs Arch. Pharmacol. 279: 207-210 (1973).
152. Seydel, J. K., and Wassermann, O. NMR-studies on the molecular basis of drug-induced phospholipidosis. II. Interaction between several amphiphilic drugs and phospholipids. *Biochim. Pharmacol.* 25: 2357-2364 (1976).
 153. Ohkuma, S., and Poole, B. Fluorescence probe measurement of the intralysosomal pH in living cells and the perturbation of pH by various agents. *Proc. Natl. Acad. Sci. (U.S.)* 75: 3327-3331 (1978).
 154. Kasama, K., Yoshida, K., Takeda, S., Akeda, S., and Kawai, K. Bis-(monoacylglyceryl)phosphate and acyl phosphatidylglycerol isolated from human livers of lipidosis induced by 4,4'-diethylaminoethoxyhexestrol. *Lipids* 9: 235-243 (1974).
 155. Wherrett, J. R., and Huterer, S. Bis-(monoacylglyceryl)-phosphate of rat and human liver: fatty acid composition and NMR spectroscopy. *Lipids* 8: 531-533 (1973).
 156. Yamamoto, A., Adachi, S., Ishikawa, K., Yokomura, T., Kitani, T., Nasu, T., Imoto, T., and Nishikawa, M. Studies on drug-induced lipidosis. III. Lipid composition of the liver and some other tissues in clinical cases of "Niemann-Pick-like syndrome" induced by 4,4'-diethylaminoethoxyhexestrol. *J. Biochem.* 70: 775-784 (1971).
 157. Blohm, T. R. Drug-induced lysosomal lipidosis: biochemical interpretations. *Pharmacol. Rev.* 30: 593-603 (1979).
 158. Brindley, D. N., and Bowley, M. Drugs affecting the synthesis of glycerides and phospholipids in rat liver. *Biochem. J.* 148: 461-469 (1975).
 159. Bowley, M., Cooling, J., Burditt, S. L., and Brindley, D. N. The effects of amphiphilic cationic drugs and inorganic cations on the activity of phosphatidate phosphohydrolase. *Biochem. J.* 165: 447-454 (1977).
 160. Colbeau, A., Cuault, F., and Vignais, P. M. Characterization and subcellular localization of lipase activities in rat liver cell. Comparison with phospholipase A. *Biochimie* 56: 275-288 (1974).
 161. Mellors, A., and Tappel, A. L. Hydrolysis of phospholipids by a lysosomal enzyme. *J. Lipid Res.* 8: 479-485 (1967).
 162. Gräbner, R. Action of chlorpheniramine on the hydrolysis of phosphatidylcholine by phospholipase A₂. *Exptl. Pathol.* 13: 68-77 (1977).
 163. Hostetler, K. Y., and Matsuzawa, Y. Studies on the mechanism of drug-induced lipidosis. Cationic amphiphilic drug inhibition of lysosomal phospholipases A and C. *Biochem. Pharmacol.* 30: 1121-1126 (1981).
 164. Kunze, H., Nahas, N., Traynor, J. R., and Wurl, M. Effects of local anesthetics on phospholipases. *Biochim. Biophys. Acta* 441: 93-102 (1976).
 165. Hostetler, K. Y., and Hall, L. B. Inhibition of kidney lysosomal phospholipases A and C by aminoglycoside antibiotics: possible mechanism of aminoglycoside toxicity. *Proc. Natl. Acad. Sci. (U.S.)* 79: 1663-1667 (1982).
 166. Matsuzawa, Y., and Hostetler, K. Y. Inhibition of lysosomal phospholipase A and phospholipase C by chloroquine and 4,4'-bis(diethylaminoethoxy)- α,β -diethyldiphenylethane. *J. Biol. Chem.* 255: 5190-5194 (1980).
 167. Stremmel, W., and Debuch, H. Bis(monoacylglycerol)phosphate—a marker lipid of secondary lysosomes. *Hoppe-Seylers Z. Physiol. Chem.* 357: 803-810 (1976).
 168. Tjong, H. B., Lepthin, J., and Debuch, H. Lysosomal phospholipids from rat liver after treatment with different drugs. *Hoppe-Seylers Z. Physiol. Chem.* 359: 63-67 (1978).
 169. Bleistein, J., Debuch, H., and Gunawan, J. Metabolism of phosphatidylglycerol by liver lysosomes of chloroquine-pretreated rats. *Hoppe-Seylers Z. Physiol. Chem.* 361: 1445-1448 (1980).
 170. Frentzen-Bertrams, M., and Debuch, H. Production of bis(monoacylglycerol)phosphate from phosphatidylglycerol in isolated lysosomes of chloroquine-pretreated rats. *Hoppe-Seylers Z. Physiol. Chem.* 362: 1229-1236 (1981).
 171. Wherrett, J. R., and Huterer, S. Enrichment of bis-(monoacylglyceryl)phosphate in lysosomes from rat liver. *J. Biol. Chem.* 247: 4114-4120 (1972).
 172. Hendelman, W. J. A morphologic study of the effects of LSD on neurons in cultures of cerebellum. *J. Neuropathol. Exptl. Neurol.* 31: 411-432 (1972).
 173. Trout, J. J., Stauber, W. T., and Schottelius, B. A. Increased autophagy in chloroquine-treated tonic and phasic muscles: an alternative view. *Tissue Cell* 13: 393-401 (1981).
 174. Gerber, N., and Arnold, K. The effect of diphenyl-piperazine compounds and other agents on diphenhydantoin, zoxazolamine and hexobarbital metabolism. *J. Pharmacol. Exptl. Therap.* 164: 232-238 (1968).
 175. Leeson, G. A., Biedenbach, S. A., Chan, K. Y., Gibson, J. P., and Wright, G. J. Decrease in the activity of the drug-metabolizing enzymes of rat liver following the administration of tilorone hydrochloride. *Drug Metab. Dispos.* 4: 232-238 (1976).
 176. Hers, H. G., and Van Hoff, F. (Eds.). *Lysosomes and Storage Diseases*. Academic Press, New York, 1973.
 177. Buchheim, W., Drenckhahn, D., and Lüllmann-Rauch, R. Freeze-fracture studies of cytoplasmic inclusions occurring in experimental lipidosis as induced by amphiphilic cationic drugs. *Biochim. Biophys. Acta* 575: 71-80 (1979).
 178. Lüllmann-Rauch, R. Retinal lipidosis in albino rats treated with chlorpheniramine and with tricyclic antidepressants. *Acta Neuropathol. (Berlin)* 35: 55-67 (1976).
 179. Howell, J. I., Fisher, D., Goodall, A. H., Verrinder, M., and Lucy, J. A. Interactions of membrane phospholipids with fusogenic lipids. *Biochim. Biophys. Acta* 332: 1-10 (1973).
 180. Junger, E., and Reinauer, H. Liquid crystalline phases of hydrated phosphatidylethanolamine. *Biochim. Biophys. Acta* 183: 304-308 (1969).
 181. Cain, H., and Kraus, B. Lysosomal pigmentary deposits in the kidney of old Wistar rats. *Virchows Arch. (Cell Pathol.)* 10: 322-338 (1972).
 182. Feldman, S., Wang, M. Y., and Kaloyanides, G. J. Aminoglycosides induce a phospholipidosis in the renal cortex of the rat: An early manifestation of nephrotoxicity. *J. Pharmacol. Exptl. Therap.* 220: 514-520 (1982).
 183. Kosek, J. C., Mazze, R. I., and Cousins, M. J. Nephrotoxicity of gentamycin. *Lab. Invest.* 30: 48-57 (1974).
 184. Luft, F. C., Yum, M. N., Walker, P. D., and Kleit, S. A. Gentamicin gradient patterns and morphological changes in human kidneys. *Nephron* 18: 167-174 (1977).
 185. Fowler, B. A., and Brooks, R. E. Effects of the herbicide paraquat on the ultrastructure of mouse kidney. *Am. J. Pathol.* 63: 505-512 (1971).
 186. France, R., Gray, M. E., Stone, W. J., and Swift, L. L. Intracellular granules of the renal medulla in a case of potassium depletion due to renal potassium wasting. *Am. J. Pathol.* 91: 299-312 (1978).
 187. Morré, J., Kartenbeck, J., and Frane, W. W. Membrane flow and interconversions among endomembranes. *Biochim. Biophys. Acta* 559: 71-152 (1979).
 188. Hruban, Z., Tavoloni, N., Reed, J. S., and Boyer, J. L. Ultrastructural changes during cholestasis induced by chlorpromazine in the isolated perfused rat liver. *Virchows Arch. (Cell Pathol.)* 26: 289-305 (1978).
 189. Kvarstein, B., and Stormorken, H. Influence of acetylsalicylic acid, butazolidine, colchicine, hydrocortisone, chlorpromazine and imipramine on the phagocytosis of polystyrene latex particles by human leucocytes. *Biochem. Pharmacol.* 20: 119-124 (1971).
 190. Keeffe, E. B., Blankenship, N. M., and Scharschmidt, B. F. Alteration of rat liver plasma membrane fluidity and ATPase activity by chlorpromazine hydrochloride and its metabolites. *Gastroenterology* 79: 222-231 (1980).
 191. Mizuno, G. R., Chapman, C. J., Chipault, J. R., and Pfeiffer, D. R. Lipid composition and (Na⁺ + K⁺)-ATPase activity in rat lens during triparanol-induced cataract formation. *Biochim. Biophys. Acta* 664: 1-12 (1981).
 192. Williams, P. D., Holohan, P. D., and Ross, C. R. Gentamycin nephrotoxicity. I. Acute biochemical correlates in rats. *Toxicol. Appl. Pharmacol.* 61: 234-242 (1981).
 193. Bermejo, J., Fernandez, P., Tato, F., and Belmonte, A. Desmethylinipramine induced changes in liposomal membranes. *Res. Commun. Chem. Pathol. Pharmacol.* 8: 101-113 (1974).

194. Tsao, S. C., Iga, T., Sugiyama, Y., and Hanano, M. Effect of chlorpromazine on isolated rat hepatocytes. *Biochem. Pharmacol.* 31: 491-497 (1982).
195. Cater, B. R., Chapman, D., Hawes, S. M., and Saville, J. Lipid phase transitions and drug interactions. *Biochim. Biophys. Acta* 363: 54-69 (1974).
196. Jain, M. K., Wu, N. Y., and Wray, L. V. Drug-induced phase change in bilayer as possible model of action of membrane expanding drugs. *Nature* 255: 494-495 (1975).
197. Rothman, J. E. The molecular basis of mesomorphic phase transitions in phospholipid systems. *J. Theor. Biol.* 38: 1-16 (1973).
198. Watts, S. D. M., and Atkins, A. M. Effects of the schistosomicide, 1,7-bis(*p*-aminophenoxy)heptane (153C51) on lysosomes and membrane stability. *Biochem. Pharmacol.* 28: 2579-2584 (1979).
199. Perel, J. M. Pharmacokinetics of therapeutic and toxic reactions: II. Tricyclic antidepressants. *Neurotoxicology* 1: 157-161 (1977).
200. Anzil, A. P., Herrlinger, H., and Blinzinger, K. Ultrastructural observation on the cytoplasmic inclusions of nervous tissues and peripheral myopathy induced in suckling mice by chlorphen-termine administration. *Neurotoxicology* 1: 485-491 (1977).
201. Drenckhahn, D., and Lüllmann-Rauch, R. Lens opacities associated with lipidosis-like ultrastructural alterations in rats treated with chloroquine, chlorphen-termine or iprindole. *Exp. Eye Res.* 24: 621-632 (1977).
202. Abraham, R., Hendy, R., and Grasso, P. Formation of myeloid bodies in rat liver lysosomes after chloroquine administration. *Exptl. Mol. Pathol.* 9: 212-229 (1968).
203. Miller, D. K., and Lenard, J. Antihistaminics, local anesthetics, and other amines as antiviral agents. *Proc. Natl. Acad. Sci. (U.S.)* 78: 3605-3609 (1961).
204. Pearson, R. D., Manian, A. A., Marcus, J. L., Hall, D., and Hewlett, E. L. Lethal effect of phenothiazine neuroleptics on the pathogenic protozoan, *Leishmania donovani*. *Science* 217: 369-371 (1982).
205. Shikata, T., Kanetaka, T., Endo, Y., and Nagashima, K. Drug-induced generalized phospholipidosis. *Acta Pathol. Japan* 22: 517-531 (1972).
206. Dukes, M. N. G. (Ed.) *Side Effects of Drugs Annual 3*. Excerpta Medica, Amsterdam, 1977.
207. Mielke, H., Seiler, K. U., Stumpf, U., and Wassermann, O. Über eine Beziehung zwischen dem Serotoninstoffwechsel und der pulmonalen Hypertonie bei Ratten nach Gabe verschiedener Anorektika. *Z. Kardiol.* 62: 1090-1098 (1973).
208. Angevine, L. S., and Mehendale, H. M. Effect of chlorphen-termine on the pulmonary disposition of 5-hydroxytryptamine in the isolated perfused rabbit lung. *Am. Rev. Resp. Dis.* 122: 891-898 (1980).
209. Flodh, H., Magnusson, G., and Magnusson, O. Pulmonary foam cells in rats of different age. *Z. Versuchstierkd.* 16: 299-312 (1974).
210. Aubert-Tulkens, G., Van Hoff, F., and Tulkens, P. Gentamicin-induced lysosomal phospholipidosis in cultured rat fibroblasts. *Lab. Invest.* 40: 481-491 (1979).
211. Unsicker, K., Allan, I. J., and Newgreen, D. F. Extraneuronal effects of 6-hydroxydopamine and extraneuronal uptake of noradrenaline. *In vivo* and *in vitro* studies on adrenocortical cells of lizards and rats. *Cell Tissue Res.* 173: 45-69 (1976).
212. Gray, J. E., Purmalis, A., Purmalis, B., and Mathews, J. Ultrastructural studies of the hepatic changes brought about by clindamycin and erythromycin in animals. *Toxicol. Appl. Pharmacol.* 19: 217-233 (1971).
213. Loker, E., Scallen, T. J., and Diettert, S. E. Phenobarbital-induced proliferation of smooth endoplasmic reticulum after administration of triparanol. *Anat. Rec.* 168: 221-225 (1970).
214. Svendsen, O. Effect of phenobarbital on chlorphen-termine induced alveolar foam cells in rats. *Toxicol. Appl. Pharmacol.* 40: 171-173 (1977).
215. Kim, S. U. Effects of the cholesterol biosynthesis inhibitor AY 9944 on organotypic cultures of mouse spinal cord. Retarded myelinogenesis and induction of cytoplasmic inclusions. *Lab. Invest.* 32: 720-728 (1975).
216. Rosenthal, A. R., Kolb, H., Bergsma, D., Huxsoll, D., and Hopkins, J. L. Chloroquine retinopathy in the rhesus monkey. *Invest. Ophthalmol. Vis. Sci.* 17: 1158-1175 (1978).
217. Serabjit-Singh, C. J., Wolf, C. R., Philpot, R. M., and Plopper, C. G. Cytochrome P-450: localization in rabbit lung. *Science* 207: 1467-1470 (1980).
218. Redding, R. A., Doubilas, W. H. J., and Stein, M. Thyroid hormone influence upon lung surfactant metabolism. *Science* 175: 994-996 (1972).
219. Hook, G. E. R., and DiAugustine, R. P. Secretory cells of the peripheral pulmonary epithelium as targets for toxic agents. *Environ. Health Perspect.* 16: 147-156 (1976).
220. Ashford, A., and Ross, J. W. Toxicity of depressant and antidepressant drugs in hyperthyroid mice. *Brit. Med. J.* 2: 217-218 (1968).
221. Philpot, R. M., Anderson, M. W., and Eling, T. E. Uptake, accumulation, and metabolism of chemicals by the lung. In: *Metabolic Function of the Lung*. (Y. S. Bakhle and J. R. Vane, Eds.), Marcel Dekker, New York, 1977, pp. 123-171.
222. Lüllmann-Rauch, R., and Pietschmann, N. Lipidosis-like cellular alterations in lymphatic tissues of chlorphen-termine-treated animals. *Virchows Arch. (Cell Pathol.)* 15: 295-308 (1974).
223. Bingham, E., Barkley, W., Zerwas, M., Stemmer, K., and Taylor, P. Responses of alveolar macrophages to metals. I. Inhalation of lead and nickel. *Arch. Environ. Health* 25: 406-414 (1972).
224. Deodhar, S. D., and Bhagwat, A. G. Desquamative-interstitial pneumonia-like syndrome in rabbits. *Arch. Pathol.* 84: 54-58 (1967).
225. Brewer, D. B., Heath, D., and Asquith, P. Electron microscopy of desquamative interstitial pneumonia. *J. Pathol.* 97: 317-323 (1969).
226. Corrin, B., and Price, A. B. Electron microscopic studies in desquamative interstitial pneumonia associated with asbestos. *Thorax* 27: 324-331 (1972).
227. Farr, G. H., Harley, R. A., and Hennigar, G. R. Desquamative interstitial pneumonia. An electron microscopic study. *Am. J. Pathol.* 60: 347-370 (1970).
228. Heath, D. Desquamative interstitial pneumonia. *Thorax* 23: 330 (1968).
229. Rhodes, M. L. Desquamative interstitial pneumonia. New ultrastructural findings. *Am. Rev. Respir. Dis.* 108: 950-954 (1973).
230. Shortland, J. R., Drake, C. S., and Crane, W. A. J. Electron microscopy of desquamative interstitial pneumonia. *Thorax* 24: 192-208 (1969).
231. Bhagwat, A. G., Wentworth, P., and Conen, P. E. Observations on the relationship of desquamative interstitial pneumonia and pulmonary alveolar proteinosis in childhood: a pathologic and experimental study. *Chest* 58: 326-332 (1970).
232. Heppleston, A. G., Wright, N. A., and Stewart, J. A. Experimental alveolar lipo-proteinosis following the inhalation of silica. *J. Pathol.* 101: 293-307 (1970).
233. Kuhn, C., Györkey, G., Levine, B. E., and Ramirez, R. J. Pulmonary alveolar proteinosis. A study using enzyme, histochemistry, electron microscopy and surface tension measurements. *Lab. Invest.* 15: 492-509 (1966).
234. Rosen, S. H., Castleman, B., and Liebow, A. A. Pulmonary alveolar proteinosis. *N. Engl. J. Med.* 258: 1123-1142 (1958).
235. Davidson, J. M., and Macleod, W. M. Pulmonary alveolar proteinosis. *Brit. J. Dis. Chest* 63: 13-28 (1969).
236. Divertie, M. B., Brown, A. L., and Harrison, E. G. Pulmonary alveolar proteinosis. Two cases studied by electron microscopy. *Am. J. Med.* 40: 351-359 (1966).
237. Schober, R., Kosek, J. C., and Bensch, K. G. The origin of the membranaceous intralveolar material in pulmonary alveolar proteinosis. *Lab. Invest.* 30: 388-389 (1974).
238. Lamberty, J., Hoffmann, E., and Pizzolato, P. The ultrastructure of acute silicosis in humans. *Am. J. Pathol.* 70: 34a-35a (1973).
239. Hoffmann, E. O., Lamberty, J., Pizzolato, P., and Coover, J. The ultrastructure of acute silicosis. *Arch. Pathol.* 96: 104-107

- (1973).
240. Costello, J. F., Moriarty, D. C., Branthwaite, M. A., Turner-Warwick, M., and Corrin, B. Diagnosis and management of alveolar proteinosis: the role of electron microscopy. *Thorax* 30: 121-132 (1975).
 241. Gleiser, C. A., Bay, W., Dukes, T. W., Brown, R. S., Read, W. K., and Pierce, K. R. Study on chloroquine toxicity and a drug-induced cerebrosplinal lipodystrophy in swine. *Am. J. Pathol.* 53: 27-45 (1968).
 242. Itoh, S., and Tsukada, Y. Clinico-pathological and electron microscopical studies on a coronary dilating agent: 4,4'-diethyl-aminoethoxyhexestrol-induced liver injuries. *Acta Hepatogastroenterol.* 20: 204-215 (1973).
 243. Bockhardt, H., Drenckhahn, D., and Lüllmann-Rauch, R. Generalisierte Lipidose bei der Ratte nach Behandlung mit Amiodaron und Zimelidin. *Verh. Anat. Ges.* 72: 417-418 (1978).
 244. Theiss, E., Hummler, H., Lengsfeld, H., Staiger, G. R., and Tranzer, J. P. Lipidspeicherung bei Versuchstieren nach Verabreichung trizyklischer Amine. *Schweiz. Med. Wochenschr.* 103: 424 (1973).
 245. Brown, E. A. B. The localization, metabolism and effects of drugs and toxicants in lung. *Drug Metab. Rev.* 3: 33-87 (1974).
 246. Ryrfeldt, A. The distribution, elimination, and biotransformation of ¹⁴C-Cloforex in the mouse and rat. *Acta Pharmacol. Toxicol. (Copenh.)* 28: 391-405 (1970).
 247. Mazue, G., Berthe, J., Newmann, A. J., and Brunaud, M. A toxicologic evaluation of ethyl fluclozepat (CM 6912). *Int. J. Clin. Pharmacol. Therap. Toxicol.* 19: 453-472 (1981).
 248. Kacaw, S., and Narbaitz, R. A comparative ultrastructural and biochemical study between the effects of chlorphentermine and phentermine on rat lung. *Exptl. Mol. Pathol.* 27: 106-120 (1977).
 249. Franken, G., Lüllmann, H., and Siegfriedt, A. The occurrence of huge cells in pulmonary alveoli of rats treated by an anorexic drug. *Arzneim. Forsch.* 20: 417 (1970).
 250. Heath, D., Smith, P., and Hasleton, P. Effect of chlorphentermine on the rat lung. *Thorax* 28: 551-558 (1973).
 251. Eling, T. E., Pickett, R. D., Orton, T. C., and Anderson, M. W. A study of the dynamics of imipramine accumulation in the isolated perfused rabbit lung. *Drug Metab. Dispos.* 3: 389-399 (1975).
 252. D'Amico, D. J., Kenyon, K. R., and Ruskin, J. N. Amiodarone keratopathy. Drug-induced lipid storage disease. *Arch. Ophthalmol.* 99: 257-261 (1981).
 253. Yates, R. D., Chen, I. L., and Mascorro, J. A. Some morphological effects of 20,25-diazacholesterol (SC-12937) on adrenocortical cells of the Syrian hamster. *Tex. Rep. Biol. Med.* 26: 241-248 (1968).
 254. Batty, H. K., and Millhouse, O. E. Ultrastructure of the Gunn rat substantia nigra I. Cytoplasmic changes. *Acta Neuropathol.* 35: 93-107 (1976).
 255. Schutta, H. S., and Johnson, L. Bilirubin encephalopathy in the Gunn rat. A fine structural study of the cerebellar cortex. *J. Neuropathol. Exptl. Neurol.* 26: 377-396 (1967).
 256. Kacaw, S., and Narbaitz, R. Early metabolic alterations in pulmonary tissue following administration of toxic agents. *Fed. Proc.* 37: 2489-2495 (1978).
 257. Hassenpflug, J. Further studies on the structure-activity relationship of drugs inducing lipodosis-like ultrastructural alterations. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 293 (Suppl): R57 (1976).
 258. Grunnet, M. L. Effects of nortriptyline and other psychotropic drugs on neurons and glia *in vitro*: an ultrastructural study. *Neurotoxicology* 1: 163-170 (1977).
 259. Grant, W. M. Toxicology of the eye. Charles C Thomas, Springfield, IL, 1974.
 260. Merrill, T. G., Payne, B. J., and Tousimis, A. J. Cytoplasmic granules in lymphocytes from rats dosed with SK&F 14336-D. In: *Proceedings 29th Annual Meeting, Electron Microscopy Society of America* (C. J. Arceneaux, Ed.), Claitor's Publishing Division, Baton Rouge, LA, 1971, pp. 556-557.
 261. Weiss, J. N., Weinberg, R. S., and Regelson, W. Keratopathy after oral administration of tilorone hydrochloride. *Am. J. Ophthalmol.* 89: 46-53 (1980).
 262. Weiss, J. N., Ochs, A. L., Abedi, S., and Selhorst, J. B. Retinopathy after tilorone hydrochloride. *Am. J. Ophthalmol.* 90: 846-853 (1980).
 263. Drenckhahn, D., and Lüllmann-Rauch, R. Drug-induced experimental lipodosis in the nervous system. *Neuroscience* 4: 697-712 (1979).
 264. Thompson, S. W., Antonchak, B., Matthews, J. R., and Katz, R. Chloroquine induced retinal gangliopathy. *Bull. Soc. Pharmacol. Environ. Pathol.* 3: 2-24 (1975).
 265. Fedorko, M. E. Effect of chloroquine on morphology of leukocytes and pancreatic exocrine cells from the rat. *Lab. Invest.* 18: 27-37 (1968).
 266. Drenckhahn, D., and Lüllmann-Rauch, R. Experimental myopathy induced by amphiphilic cationic compounds including several psychotropic drugs. *Neuroscience* 4: 549-562 (1979).
 267. Rawlins, F. A., and Uzman, B. G. Effect of AY-9944, a cholesterol biosynthesis inhibitor on peripheral nerve myelination. *Lab. Invest.* 23: 184-189 (1970).
 268. Abraham, R., and Hendy, R. Irreversible lysosomal damage induced by chloroquine in the retinae of pigmented and albino rats. *Exptl. Mol. Pathol.* 12: 185-200 (1970).
 269. Read, W. K., and Bay, W. W. Basic cellular lesions in chloroquine toxicity. *Lab. Invest.* 24: 246-259 (1971).
 270. Tousimis, A. J., and Barron, C. N. Chlorpromazine and the eye of the dog. *Exptl. Mol. Pathol.* 13: 89-110 (1970).
 271. Wong, T.-W., and Hruban, Z. Testicular degeneration and necrosis induced by chlorcyclizine. *Lab. Invest.* 26: 278-299 (1972).
 272. Houghton, D. C., Hartnett, M., Campbell-Boswell, M., Porter, G., and Bennett, W. A light and electron microscopic analysis of gentamycin nephrotoxicity in rats. *Am. J. Pathol.* 82: 589-612 (1976).
 273. Lhermitte, F., Fardeau, M., Chedru, F., and Mallecourt, J. Polyneuropathy after perhexiline maleate therapy. *Brit. Med. J.* 1: 1256 (1976).
 274. Mussini, J. M., Hauw, J. J., and Escourolle, R. Étude en microscopie électronique des lésions nerveuses, musculaires et cutanées déterminées par le perhexiline. *Acta Neuropathol.* 38: 53-59 (1977).
 275. Nick, J., Dudognon, P., Escourolle, R., Bakouche, P., Nicolle, M. H., Regnier, A., Hauw, J. J., Ermidou, S., Pollet, S., Baumann, N., Singlas, E., and Levy, J. Manifestations neurologiques en rapport avec le traitement par le maleate de perhexiline. *Rev. Neurol. (Paris)* 134: 103-114 (1978).
 276. Chen, I. L., and Yates, R. O. An ultrastructural study of opaque cytoplasmic inclusions induced by triparanol treatment. *Am. J. Anat.* 121: 705-726 (1976).
 277. Diener, R. M., and Hsu, B. Y. D. Effects of certain basic phenolic ethers on the rat fetus. *Toxicol. Appl. Pharmacol.* 10: 565-567 (1967).
 278. Schutta, H. S., and Neville, H. E. Effects of cholesterol synthesis inhibitors on the nervous system. *Lab. Invest.* 19: 487-493 (1968).
 279. Yates, R. D. The effects of triparanol on adrenocortical cells of the zona fasciculata of Syrian hamsters. *Z. Zellforsch.* 71: 41-52 (1966).
 280. James-Kracke, M. R., Sloane, B. F., Schuman, H., and Somlyo, A. P. Lysosomal composition in cultured vascular smooth muscle cells: Electron probe analysis. *Proc. Natl. Acad. Sci. (U.S.)* 76: 6461-6465 (1979).
 281. Dubois-Dalcq, J., Buyse, M., DeBruyne, J., Van Tilghen, H., and Thiry, L. Ultrastructural study of the action of aminopiperazines on transformed cell lines and on nervous system tissue culture. *Exptl. Cell Res.* 77: 303-311 (1973).
 282. Goodwin, L. G., Richards, W. H. G., and Udall, V. The toxicity of diaminodiphenoxyalkanes. *Brit. J. Pharmacol.* 12: 468-474 (1957).
 283. Drenckhahn, D., Kleine, L., and Lüllmann-Rauch, R. Lysosomal alterations in cultured macrophages exposed to anorexiogenic and psychotropic drugs. *Lab. Invest.* 35: 116-123 (1976).
 284. Cowger, M. L. Mechanism of bilirubin toxicity on tissue culture

- cells. Factors that affect toxicity, reversibility by albumin, and comparison with other respiratory poisons and surfactants. *Biochem. Med.* 5: 1-16 (1971).
285. Silberberg, D. H., and Schutta, H. S. The effects of unconjugated bilirubin and related pigments on cultures of rat cerebellum. *J. Neuropathol. Exptl. Neurol.* 26: 576-583 (1967).
286. Alexander, N. J. Ultrastructural changes in rat epididymis after vasectomy. *Z. Zellforsch.* 136: 177-182 (1973).
287. Liveson, J. A., Gardner, J., and Bornstein, M. B. Tissue culture studies of possible uremic neurotoxins: Myoinositol. *Kidney Int.* 12: 131-136 (1977).
288. West, N. R. Response of rat dorsal root and superior cervical ganglion cultures to congeners of chlorpromazine. *Proc. Fourth Ann. Meet. Soc. Neuroscience* 4: 474 (1974).
289. Glassy, M. C., and Ferrone, S. Ultrastructural alterations in human lymphoblastoid B cell lines treated with tunicamycin. *Am. J. Pathol.* 103: 1-9 (1981).
290. Fedorko, M. E., Hirsch, J. G., and Cohn, Z. A. Autophagic vacuoles produced *in vitro*. I. Studies on cultured macrophages exposed to chloroquine. *J. Cell Biol.* 38: 377-391 (1968).
291. Tischner, K. Effects of chloroquine on neurons of long-term cultures of peripheral and central nervous system. *Acta Neuropathol. (Berlin)* 28: 233-242 (1974).
292. Warhurst, D. C., and Hockley, D. J. Mode of action of chloroquine on *Plasmodium berghei* and *P. cynomolgi*. *Nature* 214: 935-936 (1967).
293. Brosnan, C. F., Bunge, M. B., and Murray, M. R. The response of lysosomes in cultured neurons to chlorpromazine. *J. Neuropathol. Exptl. Neurol.* 29: 337-353 (1970).
294. Lageron, A., Scotto, J., and Gautier, M. Effects of pexid on liver cell cultures. Ultrastructural and histochemical studies. *Eur. J. Clin. Pharmacol.* 19: 417-421 (1981).
295. Le Gall, J. Y., Guillouzo, A., Glaise, D., Deugnier, Y., Messner, M., and Bourel, M. Perhexiline maleate toxicity on human liver cell lines. *Gut* 21: 977-984 (1980).
296. Dugognon, P., Hauw, J. J., De Baeque, C., Escourolle, R., Derrida, J. P., and Nick, J. Neuropathie à l'amiodarone: lipodose medicamentuse. *Nouv. Presse Med.* 8: 1766-1767 (1979).
297. Buffalo, W. J., Johnson, A. W., and Sandifer, M. G. Total dosage of chlorpromazine and ocular opacities. *Am. J. Psychiatry* 124: 250-251 (1976).
298. Wellwood, J. M., Simpson, P. M., Tighe, J. R., and Thompson, A. E. Evidence of gentamycin nephrotoxicity in patients with renal allografts. *Brit. Med. J.* 3: 278-281 (1975).
299. McKeown, C. A., Swartz, M., Blom, J., and Maggiano, J. M. Tamoxifen retinopathy. *Brit. J. Ophthalmol.* 65: 177-179 (1981).
300. Kirby, T. S. Cataracts produced by triparanol (MER/29). *Trans. Am. Ophthalmol. Soc.* 65: 493-543 (1967).
301. McPherson, J. R., and Shorter, R. G. Intestinal lesions associated with triparanol. A clinical and experimental study. *Am. J. Dig. Dis.* 10: 1024-1033 (1965).
302. Hruban, Z. Pulmonary changes induced by amphiphilic drugs. *Environ. Health Perspect.* 16: 111-118 (1976).
303. Lüllmann, H., and Lüllmann-Rauch, R. Tamoxifen-induced generalized lipidosis in rats subchronically treated with high doses. *Toxicol. Appl. Pharmacol.* 61: 138-146 (1981).
304. Thelmo, W. L., and Levine, S. Renal lesions induced by tilorone and an analog. *Am. J. Pathol.* 91: 355-360 (1978).
305. Marchlinski, F. E., Gansler, T. S., Waxman, H. L., and Josephson, M. E. Amiodarone pulmonary toxicity. *Ann. Int. Med.* 97: 839-845 (1982).
306. Geffer, W. B., Epstein, D. M., Pietra, G. G., and Miller, W. T. Lung disease caused by amiodarone, a new antiarrhythmic agent. *Radiology* 147: 339-344 (1983).
307. Pieterse, A. S., Rowland, R., and Dunn, D. Perhexiline maleate induced cirrhosis. *Pathology* 15: 201-203 (1983).
308. Vinding, T., and Nielsen, N. V. Retinopathy caused by treatment with Tamoxifen in low dosage. *Acta Ophthalmol.* 61: 45-50 (1983).
309. Albert, C., and Lüllmann-Rauch, R. Ultrastructural alterations in peripheral nerve trunks of rats subchronically treated with chlorphentermine or perhexiline. *Drug Res.* 33: 125-127 (1983).
310. Angevine, L. S., and Mehendale, H. M. Chlorphentermine uptake by isolated perfused rabbit lung. *Toxicol. Appl. Pharmacol.* 52: 336-346 (1980).
311. Drenckhahn, D., Jacobi, B., and Lüllmann-Rauch, R. Corneal lipidosis in rats treated with amphiphilic cationic drugs. *Arzneim. Forsch.* 33(1): 827-831 (1983).
312. Ferin, J. Alveolar macrophage mediated pulmonary clearance suppressed by drug-induced phospholipidosis. *Exp. Lung. Res.* 4: 1-10 (1982).
313. Fredman, P., Klinghardt, G. W., Nilsson, O., and Svennerholm, L. Lipid accumulation in liver, spleen, lungs and kidneys of miniature pigs after chloroquine treatment. *Biochem. J.* 201: 581-588 (1982).
314. Freise, J., Magerstedt, P., and Schmid, K. Inhibition of phospholipase A₂ by gabexate mesilate, camostat and aprotinine. *Enzyme* 30: 209-212. (1983).
315. Gairola, C., Matulionis, D. H., and Reasor, M. J. Chlorphentermine-induced alterations in the lungs of vitamin E-deficient and supplemented rats. Biochemical and morphometric analysis of the pulmonary response. *Exptl. Mol. Pathol.* 38: 368-379 (1983).
316. Hirst, L. W., Sanborn, G., Green, R., Miller, N. R., and Heath, W. D. Amodiaquine ocular changes. *Arch. Ophthalmol.* 100: 1300-1304 (1982).
317. Hostetler, K. Y., and Richman, D. D. Studies on the mechanism of phospholipid storage induced by amantadine and chloroquine in Madin Darby canine kidney cells. *Biochem. Pharmacol.* 31: 3795-3799 (1982).
318. Houghton, D. C., Plamp, C. E., DeFehr, J. M., Bennett, W. H., Porter, G., and Gilbert, D. Gentamicin and tobramycin nephrotoxicity. A morphologic and functional comparison in the rat. *Am. J. Pathol.* 93: 137-152 (1978).
319. Kacew, S. Alterations in newborn and adult rat lung morphology and phospholipid levels after chlorcyclizine and chlorphentermine treatment. *Toxicol. Appl. Pharmacol.* 65: 100-108 (1982).
320. Kunze, H., Hesse, B., and Bohn, E. Effects of antimalarial drugs on several rat-liver lysosomal enzymes involved in phosphatidylethanolamine catabolism. *Biochim. Biophys. Acta* 713: 112-117 (1982).
321. Lageron, A., and Saffroy, M. Morphological histochemical biochemical changes induced by homochlorcyclizine in rat. *Acta Histochem.* 71: 103-110 (1982).
322. Lehnert, B. E., and Ferin, J. Particle binding, phagocytosis, and plastic substrate adherence characteristics of alveolar macrophages from rats acutely treated with chlorphentermine. *J. Retic. Soc.* 33: 293-303 (1983).
323. Lockard, V. G., Watson, S. A., Siraj, M. Y., Hayes, A. W., and O'Neal, R. M. Rubratoxin B hepatotoxicity: an electron microscopic study. *Exptl. Mol. Pathol.* 34: 94-109 (1981).
324. Lüllmann-Rauch, R., and Nüssberger, L. Citalopram-induced generalized lipidosis in rats. *Acta Pharmacol. Toxicol.* 52: 161-167 (1983).
325. Lüllmann-Rauch, R. Experimentally induced lipidosis in rat retinal pigment epithelium. *Arch. Klin. Ophthalmol.* 215: 297-303 (1981).
326. Lüllmann-Rauch, R. Histochemical evidence for lysosomal storage of acid glycosaminoglycans in splenic cells of rats treated with tilorone. *Histochemistry* 76: 71-87 (1982).
327. Lüllmann-Rauch, R., and Reil, G. H. Chlorphentermine-induced lipidosis-like ultrastructural alterations in lungs and adrenal glands of several species. *Toxicol. Appl. Pharmacol.* 30: 408-421 (1974).
328. Lüllmann-Rauch, R., and Stoermer, B. Generalized lipidosis in newborn rats and guinea pigs induced during prenatal development by administration of amphiphilic drugs to pregnant animals. *Virchows Arch. (Cell Pathol.)* 39: 59-73 (1982).
329. Matulionis, D. H., Gairola, C., and Reasor, M. J. Chlorphentermine-induced alterations in the lungs of vitamin E-deficient and -supplemented rats: quantitative ultrastructural analysis of type II cell response. *Exptl. Mol. Pathol.* 39: 80-88 (1983).

330. McNamara, T. E., Goodloe, S., and Butkus, D. E. Myeloid bodies in patients without clinical Fabry's disease. *Arch. Pathol. Lab. Med.* 104: 14-16 (1980).
331. Montgomery, M. R., Furry, J. M., and Reasor, M. J. Chlorphentermine inhibits oxidative energy metabolism in rat lung slices. *Toxicol. Appl. Pharmacol.* 65: 63-68 (1982).
332. Ohmiya, Y., Angevine, L. S., and Mehendale, H. M. Effect of drug-induced phospholipidosis on pulmonary disposition of pneumophilic drugs. *Drug Metab. Dispos.* 11: 25-30 (1983).
333. Pappu, A. S., and Hauser, G. Phospholipid metabolism changes in rat tissues *in vitro* after injections of propranolol. *J. Pharmacol. Exptl. Therap.* 222: 109-115 (1982).
334. Reasor, M. J., and Heyneman, C. A. Disaturated phosphatidylcholine in the pulmonary airspaces of rats treated with chlorphentermine. *Biochem. Pharmacol.* 32: 939-941 (1983).
335. Reasor, M. J., Heyneman, C. A., and Walker, E. R. Chlorcyclizine-induced pulmonary phospholipidosis in rats. *Res. Comm. Chem. Pathol. Pharmacol.* 38: 235-246 (1982).
336. Reasor, M. J., Koshut, R. A., McNulty, M. J., and Trush, M. A. Chemiluminescence from rat alveolar macrophages following induction of phospholipidosis with chlorphentermine. *Toxicol. Appl. Pharmacol.* 52: 497-506 (1980).
337. Rees, S., Constantopoulos, G., Barranger, J. A., and Brady, R. O. Organomegaly and histopathology in an animal model of mucopolysaccharidosis induced by suramin. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 319: 262-270 (1982).
338. Tashiro, Y., Watanabe, Y., and Enomoto, Y. Experimental phospholipidosis induced by 4,4'-diethyl-aminoethoxyhexestrol. *Acta Pathol. Japan.* 33: 929-942 (1983).
339. Thoma-Laurie, D. L., Walker, E. R., and Reasor, M. Chlorphentermine-induced neonatal and maternal pulmonary phospholipidosis in rats. *Exptl. Mol. Pathol.* 38: 310-321 (1983).
340. Watanabe, M. Drug-induced lysosomal changes and nephrotoxicity in rats. *Acta Pathol. Japan.* 28: 867-889 (1978).